

# Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BBG021-07

会場:301A

時間:5月26日 16:00-16:15

## 大型有孔虫 - 微細藻共生系を促進・崩壊させる環境要因 Environmental factors that enhance or collapse foraminifer-microalgal symbiosis

藤田 和彦<sup>1\*</sup>

Kazuhiko Fujita<sup>1\*</sup>

<sup>1</sup> 琉球大学理学部物質地球科学科

<sup>1</sup> University of the Ryukyus

Algal symbionts-bearing large benthic foraminifers are known as prolific primary and carbonate producers in coral-reef associated environments at the present and geological times. Understanding the mechanism of foraminifer-microalgal symbiosis is necessary to clarify their responses to secular variations in the Earth system, paleoceanographic changes, and future global environmental changes. Here I reviewed environmental factors that enhance or collapse the foraminifer-microalgal symbiosis. Negative factors which reduce net primary production of algal symbionts and the growth of a foraminiferal host include extremely high and low temperature, low pH, stagnant water motions and high nutrient concentrations. On the other hand, positive factors which enhance net primary production and calcification are poorly known. Our new culturing results indicated that higher pCO<sub>2</sub> enhanced the calcification of symbiont-bearing hyaline taxa due to enhanced photosynthetic activity of the symbionts under high pCO<sub>2</sub> seawater. Future culturing experiments under unrealistic seawater chemistry will give new insights into the mechanisms of foraminiferal-microalgal symbiosis.