

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



BPO003-P02

会場:コンベンションホール

時間:5月26日 14:00-16:30

底生有孔虫の細胞質に由来する有機物の微好気/嫌気的環境下での長期保存 Long-term preservation of organic carbon originated from the foraminiferal cell under dysoxic/anoxic conditions

野牧 秀隆^{1*}, 小川 奈々子¹, 大河内 直彦¹, 北里 洋¹

Hidetaka Nomaki^{1*}, Nanako O. Ogawa¹, Naohiko Ohkouchi¹, Hiroshi Kitazato¹

¹ 海洋研究開発機構

¹JAMSTEC

Benthic foraminifera are major components among deep-sea benthic ecosystems and are known as major consumers of phyto-detritus. However, subsequent digestion and mineralization processes of ingested organic matters are still unclear. Here we operated a long-term incubation experiment to reveal how the ingested organic matters were degraded or mineralized by benthic foraminifera under dysoxic to anoxic conditions. Two different organic matters, algae and bacteria, were labeled with ¹³C and fed onto the surface sediments in a closed system of *in situ* feeding core. After 1.5 years, the sediments were recovered and foraminiferal specimens were isolated from the sediments. Although the ¹³C-label was found at extremely low abundances in the bulk sediments and in overlying water as dissolved inorganic carbon, the label was found at high concentration in dead foraminiferal tests particularly in algae-added cores. Benthic foraminiferal feeding affects largely to the distribution and subsequent preservation/mineralization of organic matters produced at both the oceanic surface and seafloor. It further suggests foraminiferal test serves highly isolated space for preserving organic matters from surrounding environments.

キーワード: 堆積物 水境界, 底生有孔虫, 有機物, 無機化, 埋没

Keywords: Sediment-water interface, Benthic foraminifera, organic matter, Carbon mineralization, burial