

BPT002-04

会場:201B

時間:5月25日 09:15-09:30

## 完新世中期の西太平洋熱帯域表層における海水温および海水の酸素同位体比の復元 Reconstruction of temperature and oxygen isotopes in the surface water of the Mid-Holocene tropical western Pacific

井上 麻夕里<sup>1\*</sup>, Terrence M. Quinn<sup>2</sup>, Frederick W. Taylor<sup>2</sup>, 鈴木 淳<sup>3</sup>, 石村 豊穂<sup>3</sup>, 川幡 穂高<sup>1</sup>, Hai Cheng<sup>4</sup>, R. Lawrence Edwards<sup>4</sup>

Mayuri Inoue<sup>1\*</sup>, Terrence M. Quinn<sup>2</sup>, Frederick W. Taylor<sup>2</sup>, Atsushi Suzuki<sup>3</sup>, Toyoho Ishimura<sup>3</sup>, hodaka kawahata<sup>1</sup>, Hai Cheng<sup>4</sup>, R. Lawrence Edwards<sup>4</sup>

<sup>1</sup> 東大・大気海洋研, <sup>2</sup> テキサス大学地球物理研究所, <sup>3</sup> 産総研・地質情報研究部門, <sup>4</sup> ミネソタ大学地質地球物理学科  
<sup>1</sup>AORI, the Univ. Tokyo, <sup>2</sup>UTIG, the Univ. Texas, <sup>3</sup>GSI, AIST, <sup>4</sup>Dep. Geol & Geophys, Univ Minnesota

Oxygen isotope ratios ( $d^{18}\text{O}$ ) in foraminifera from deep sea cores have been used successfully to reconstruct continental ice volume, sea level and deep-ocean temperatures throughout the last glacial-interglacial cycle. Yet there is still considerable debate regarding the relative contributions of changes in the oxygen isotopic composition of seawater in the tropical surface ocean. Obtaining accurate estimate of the past  $d^{18}\text{O}$  distribution in tropical surface waters is crucial to establishing the role of the tropical oceans in global climate change. In this study, Sr/Ca ratios and  $d^{18}\text{O}$  values in a fossil coral collected from Vanuatu with the age of 5.2 ka were measured to reconstruct past sea surface temperature (SST) and  $d^{18}\text{O}$  in seawater. XRD analysis and SEM observation revealed that skeletal material was composed of only aragonite and the age determination was precisely conducted using U-Th dating method. Then Sr/Ca and  $d^{18}\text{O}$  were analyzed with an approximate time resolution of 2 month. As a result, slightly higher mean SST and about 0.4 per mil enrichment of  $d^{18}\text{O}$  relative to modern seawater have shown from 5.2 ka coral. The result is corresponding to the previous study from Great Barrier Reef which has suggested that the temperature increase enhanced the evaporative enrichment of  $^{18}\text{O}$  in seawater.

Keywords: western Pacific, coral Sr/Ca, oxygen isotope, Holocene