

BPT022-12

会場:104

時間:5月24日 16:00-16:15

## 太古代の堆積岩に含まれるパイライトの形態とその鉄同位体局所分析 The Late Archean diversity of organisms: evidence from morphology and in situ iron isotope analyses of pyrites in variou

吉屋 一美<sup>1\*</sup>, 西澤 学<sup>2</sup>, 上野 雄一郎<sup>1</sup>, 小宮 剛<sup>3</sup>, 丸山 茂徳<sup>1</sup>, 吉田 尚弘<sup>4</sup>, 平田 岳史<sup>5</sup>

Kazumi Yoshiya<sup>1\*</sup>, Manabu Nishizawa<sup>2</sup>, Yuichiro Ueno<sup>1</sup>, Tsuyoshi Komiya<sup>3</sup>, Shigenori Maruyama<sup>1</sup>, Naohiro Yoshida<sup>4</sup>, Takafumi Hirata<sup>5</sup>

<sup>1</sup> 東京工業大学 理工学研究科, <sup>2</sup> 海洋研究開発機構, <sup>3</sup> 東京大学大学院総合文化研究科, <sup>4</sup> 東京工業大学大学院総合理工学研究科, <sup>5</sup> 京都大学理学研究科

<sup>1</sup>Tokyo Institute of Technology, <sup>2</sup>JAMSTEC, <sup>3</sup>The University of Tokyo, <sup>4</sup>Tokyo Institute of Technology, <sup>5</sup>Kyoto University

The timing of the emergence and flourish of oxygenic photosynthetic organisms is still controversial. However, it is one of the key issues of the biological evolution in early Earth. Reconstruction of redox state of the late Archean seawater is important for understanding the relationship between biological activity and oxygenation.

Ferrous and its compounds were one of the major reduced species in the anoxic Archean seawater. It is widely recognized that the iron isotopic ratio changes largely through redox reactions, especially oxygen limited environment (Beard and Johnson, 2004; Johnson et al., 2004). The iron isotopic ratio of marine sedimentary minerals is useful for understanding the ocean's redox state and iron biogeochemistry in the geological past (e.g., Johnson et al., 2008). Based on the detailed geological survey of the Fortescue Group in Redmont area, we carefully selected 44 samples, which include stromatolitic carbonate rock, sandstone, mudstone, and alternation of calcareous sandstone and mudstone. We analyzed  $\delta^{56}\text{Fe}$  value of 225 pyrite grains in these samples, and discovered an extremely large variation of  $\delta^{56}\text{Fe}$  value from -4.1 to 3.0 permil.

From the result of microscopic observation, we found the relationship between pyrite grain morphology and iron isotope ratio. Most of pyrite grains with the positive  $\delta^{56}\text{Fe}$  values show hexagonal, rectangle, and parallelogram shapes, which are consistent with crystal system of iron-oxides: hematite, magnetite, and goethite, respectively. In contrast, pyrite grains with the negative  $\delta^{56}\text{Fe}$  values show pseudo-hexagonal and irregular forms. The pseudo-hexagonal shape corresponds to a monoclinic system that is crystal system of pyrrhotite. The correlation allows the possibility to solve the origin and the formation process of each grain of pyrite. We estimate the  $\delta^{56}\text{Fe}$  value of seawater from the positive  $\delta^{56}\text{Fe}$  value pyrites in stromatolitic carbonate rock and experimental data of previous studies (Welch et al., 2003; Butler et al., 2005). Based on this estimation, we consider that the  $\delta^{56}\text{Fe}$  value of pyrite lower than estimated  $\delta^{56}\text{Fe}$  value of sulfide was formed biologically. The quite low carbon isotope ratio of organic carbon with a nadir down to -47 permil indicates the activity of aerobic or anaerobic methanotrophy. On the other hand, the lowest  $\delta^{56}\text{Fe}$  value of the pyrite grain, -4.0 permil, indicates the biological iron-reducing: dissimilatory iron-reducing microorganism (DIR) or recently found iron-dependent AOM (AOM/IR). The co-occurrence indicates the oldest evidence for activity of the iron-dependent AOM. On the other hand, the  $\delta^{56}\text{Fe}$  values of pyrite grains in the mudstone layer of upper Mingah Member have relatively wide variation from -4 to +2 permil. The positive  $\delta^{56}\text{Fe}$  values suggest the partial oxidation of iron in oxygen limited surface environment. The  $\delta^{13}\text{C}_{\text{org}}$  values are also lower than 40 permil. This suggests the presence of organic carbon from methanotrophs. Iron and carbon isotopes reveal the divergence of microorganisms in Late Archean shallow sea.

キーワード: パイライト, 酸素発生型光合成, メタン酸化, ストロマトライト, 鉄還元バクテリア

Keywords: pyrite, oxygen-producing photosynthetic bacteria, methanotrophy, stromatolite, iron reducing bacteria