

BPT023-04

会場:201B

## 時間:5月22日11:30-11:45

深海底現場での巣穴型どり:初島沖湧水サイトの例 In situ burrow casting on the deep sea: an example from the Off Hatsushima cold seep site

清家 弘治<sup>1</sup>\*, ジェンキンズ ロバート<sup>2</sup>, 渡部 裕美<sup>3</sup>, 野牧 秀隆<sup>3</sup>, 佐藤 圭<sup>2</sup> Koji Seike<sup>1</sup>\*, Robert Jenkins<sup>2</sup>, Hiromi Watanabe<sup>3</sup>, Hidetaka Nomaki<sup>3</sup>, Kei Sato<sup>2</sup>

<sup>1</sup> 港湾空港技術研究所,<sup>2</sup> 横浜国立大学環境情報学府,<sup>3</sup> 海洋研究開発機構 <sup>1</sup>Port and Airport research Institute, <sup>2</sup>Yokohama National University, <sup>3</sup>JAMSTEC

Burrows produced by marine invertebrate animals are quite important for our understanding of benthic ecology. Burrows also affect significant impact on geochemical properties of the marine sediments where their producers live, because they provide seawater into sediments. However, burrow morphology on the deep sea had been unknown to date, although numerous burrows occur on the seafloor. Here we document an experimental in situ burrow casting on the Off Hatsushima cold seep site (1173 m deep) for the first time. Casts were made with polyester resin using the ROV *Hyperdolphin* and the casting device *Anagattinger*. Anastomosing network of the small burrows and Y-shaped burrow of *Acharax johnsoni* were observed. This result indicates that complex and abundant burrow system occur under the deep seafloor. In addition, the burrows might affect subsurface geochemical properties of the sediments in the seep site. Our technique can contribute to deep sea ecology, microbiology, and geology.