

BPT024-02

会場:201B

時間:5月22日 14:30-14:45

フラスニアン / ファメニアン境界大量絶滅時の海洋還元と陸上植生の崩壊 Oceanic euxinia and destruction of land vegetation during the Frasnian/Famennian boundary mass extinction

海保 邦夫^{1*}, 谷津 進¹, 大庭 雅寛¹, ポール ゴージャン²
Kunio Kaiho^{1*}, Susumu Yatsu¹, Masahiro Oba¹, Paul Gorjan²

¹ 東北大学大学院理学研究科, ² ワシントン大学

¹Tohoku University, ²Washington University

We analyzed the organic geochemistry, $\delta^{34}\text{S}_{\text{SCAS}}$, $\delta^{34}\text{S}_{\text{sulfide}}$, $\delta^{13}\text{C}_{\text{carb}}$, and pyrite diameter from three shallow sea sections in Belgium and South China including the Frasnian/Famennian (F/F) boundary. The results showed a correlation between organic geochemical redox indicators (dibenzothiophenes and 2,3,6-trimethylarylisoprenoids), sulfur and carbon isotope ratios, pyrite diameter, and an organic geochemical indicator of land vegetation destruction (dibenzofurans). We report here these new findings on redox changes and destruction of land vegetation during the Frasnian/Famennian (F/F) boundary mass extinction. We show coincident increases in $\delta^{34}\text{S}_{\text{SCAS}}$, $\delta^{34}\text{S}_{\text{sulfide}}$, $\delta^{13}\text{C}_{\text{carb}}$, dibenzothiophenes, 2,3,6-trimethylarylisoprenoids, and dibenzofurans and a coincident decrease in pyrite diameter at or near the F/F boundary in low latitude shallow-seas. These coincidental changes show the development of oceanic euxinia and destruction of land vegetation at this time. These correlations indicate that H_2S accumulated in the ocean and input of euxinic waters to the oxic surface waters that most marine organisms inhabit, would have caused marine extinctions at the F/F boundary. Destruction of land vegetation may be related to the shallow-sea euxinia.

キーワード: 硫黄同位体比, 炭素同位体比, 堆積有機分子, フラスニアン / ファメニアン境界

Keywords: sulfur isotope, carbon isotope, sedimentary organic molecules, Frasnian-Famennian boundary