

BPT024-07

会場:201B

時間:5月22日15:45-16:00

## セノマニアン期/チューロニアン期境界における rotaliporid の絶滅と短期間海洋無酸 素事変の同時性 Short-term euxinia coinciding with rotaliporid extinctions during the Cenomanian-Turonian

transition

大庭 雅寛<sup>1\*</sup>, 海保 邦夫<sup>1</sup>, 岡部 高志<sup>1</sup>, Marcos A. Lamolda<sup>2</sup>, James D. Wright<sup>3</sup> Masahiro Oba<sup>1\*</sup>, Kunio Kaiho<sup>1</sup>, Takashi Okabe<sup>1</sup>, Marcos A. Lamolda<sup>2</sup>, James D. Wright<sup>3</sup>

1 東北大学大学院理学研究科, 2 グラナダ大学, 3 ラトガース大学

<sup>1</sup>Tohoku University, <sup>2</sup>Universidad de Granada, <sup>3</sup>Rutgers University

Oceanic anoxic event 2 (OAE2), which occurred during the Cenomanian-Turonian (C-T) transition and lasted 10<sup>6</sup> years, is characterized by a positive global carbon isotopic excursion and stepwise extinctions in marine biota. To examine temporal variations in the dissolved oxygen content of the water column, shallow-marine C-T sediments from northern Spain were analyzed for concentrations of dibenzothiophenes, which are indicators of euxinic depositional environments, and 2,3,6trimethylarylisoprenoids, which probably indicate photic-zone euxinia. The positive excursion in d<sup>13</sup>C values of carbonates is accompanied by short- ( $10^3$  to  $10^4$  years) and long-term ( $10^5$  years) increases in dibenzothiophene and 2,3,6-trimethylarylisoprenoid concentrations, suggesting that the bottom water and photic zone of the eastern marginal sea of the North Atlantic Ocean were euxinic. Two of the short-term increases in organic compound concentrations took place just after the last occurrence of the planktonic foraminifers Rotalipora greenhornensis and R. cushmani. These transient maxima indicate that the extinction of both planktonic foraminifers was due to short-term OAEs lasting  $10^3$  to  $10^4$  years.

Keywords: Cenomanian-Turonian, oceanic anoxic event, foraminiferal extinction, dibenzothiophenes, 2,3,6-trimethylarylisoprenoids, brachiopods