

BPT024-08

会場:201B

時間:5月22日 16:00-16:15

スペイン南部カラバカの白亜紀 / 古第三紀境界近傍における海洋酸化還元と陸上植生

Oceanic redox and land vegetation across the Cretaceous/Paleogene boundary in Caravaca, southern Spain

水上 拓也^{1*}, 海保邦夫¹, 大庭 雅寛¹

Takuya Mizukami^{1*}, Kaiho Kunio¹, Masahiro Oba¹

¹ 東北大学理学部地圏環境科学科

¹ Geol. and Paleontol., Tohoku Univ.

An extraterrestrial impact at the Cretaceous/Paleogene (K/P) boundary ca. 65.5 million years ago caused a mass extinction and rapid changes of surface environment on the Earth. Here we report changes of oceanic redox and land vegetation across the K/P boundary at Caravaca, southern Spain using sedimentary organic molecules. The results reveal that the basal 3-mm thin layer of the K/P boundary-clay is marked by a rapid increase in concentrations of terrestrial long-chain n-alkanes and dibenzofuran, indicating a destruction of land vegetation and an increase in supply of terrestrial organic matter into the marine environment at the K/P boundary. This layer also have a rapid increase in concentrations of dibenzothiophene, indicating a change in redox conditions from oxic to anoxic/euxinic conditions. The low-oxygen condition could have been caused by an increase in flux of terrestrial organic matter into the ocean. A rapid increase in concentrations of retene and retene/cadalene occurred in the upper part of the boundary-clay, indicating a recovery of angiosperm which withered at the K/P boundary.