

BPT026-04

会場:201B

時間:5月24日 15:00-15:15

ミャンマーにおける新第三紀のアントラコテリウム類 (哺乳綱, 偶蹄目) の進化 Evolution of the anthracotheres (Mammalia, Artiodactyla) in the Neogene of Myanmar

鏗本 武久^{1*}, タウンタイ², ジンマウンマウンテイン³, 江木直子⁴, 高井正成⁴
Takehisa Tsubamoto^{1*}, Thaung-Htike², Zin-Maung-Maung-Thein³, Naoko Egi⁴, Masanaru Takai⁴

¹ 林原生物化学研究所・古生物学研究センター, ² シュエボー大学, ³ マンダレー大学, ⁴ 京都大学・霊長類研究所
¹Hayashibara Biochemical Lab., Inc., ²Shwebo Degree Collage, ³Mandalay Univ., ⁴Primate Res. Inst., Kyoto Univ.

We report new gnatho-dental specimens of the anthracotheres (Mammalia, Artiodactyla) discovered from four Neogene localities of central Myanmar. Based on these new specimens, we recognized four species of the anthracotheres in the Neogene of central Myanmar: *Microbunodon silistrensis* and aff. *Sivameryx* sp. from the middle Miocene; and *Microbunodon milaensis* and *Merycopotamus dissimilis* from the latest Miocene to Plio-Pleistocene. This discovery extends the distribution of *Microbunodon* and *Sivameryx*-like bothriodontine from the Indian Subcontinent to Southeast Asia, indicating their broader distribution in the Neogene. Furthermore, the discovery demonstrated that *Microbunodon* survived until the Plio-Pleistocene. It also indicates that both a highly selenodont hippo-like form (*Merycopotamus*) and a bunodont and relatively primitive form (*Microbunodon*) were the last surviving anthracotheres. These two anthracotheres co-existed until the late Pliocene/early Pleistocene in Myanmar. In the Neogene of Myanmar, the anthracotheriid fauna was changed around the late Miocene. This change was probably caused by the invasion of anthracotheres from the Indian Subcontinent likely related to the major faunal turnover events in the subcontinent. Then, the later fauna persisted until the late Pliocene/early Pleistocene, when the last anthracotheres became extinct.

キーワード: ミャンマー, 新第三紀, アントラコテリウム科, 哺乳類, 化石
Keywords: Myanmar, Neogene, Anthracotheriidae, Mammalia, fossil