

HDS004-P06

会場:コンベンションホール

時間:5月27日 09:00-10:45

KH10-5 スマトラ北西沖調査航海概要—高分解能 MCS 調査— High-resolution MCS survey during KH-10-5 Leg.1 off northwest Sumatra cruise

平田 賢治^{1*}, Riza Rahardiawan², 馬場久紀³, Leonardo Seeber⁴, 亀尾桂⁵, 山本英樹⁶, 林央之⁶, 三澤文慶⁵, 安達啓太⁵, 去川寛士³, 関本岳朗⁷, 井餘田航希⁷, 富士原敏也⁸, 木下正高⁸, 徳山英一⁵, 中村恭之⁸, 荒井晃作⁹, Haryadi Permana¹⁰, Udrek¹¹, Yusuf S. Djajadihardja¹²

Kenji Hirata^{1*}, Riza Rahardiawan², Hisatochi Baba³, Leonardo Seeber⁴, Katsura Kameo⁵, Hideki Yamamoto⁶, Hiroyuki Hayashi⁶, Ayanori Misawa⁵, Keita Adachi⁵, Hiroshi Sarukawa³, Takerou Sekimoto⁷, Kohki Iyota⁷, Toshiya Fujiwara⁸, Masataka Kinoshita⁸, Hidekazu Tokuyama⁵, Yasuyuki Nakamura⁸, Kohsak Arai⁹, Haryadi Permana¹⁰, Udrek¹¹, Yusuf S. Djajadihardja¹²

¹ 気象研究所, ² インドネシア海洋地質研究所, ³ 東海大学, ⁴ ラumont・ドハティ地球観測所, ⁵ 東京大学, ⁶ マリンワークジャパン, ⁷ 高知大学, ⁸ 海洋研究開発機構, ⁹ 産業技術総合研究所, ¹⁰ インドネシア科学院, ¹¹ インドネシア応用技術評価庁, ¹² インドネシア測量地図庁

¹MRI, ²MGI, ³Tokai Univ., ⁴LDEO, ⁵Univ. of Tokyo, ⁶MWJ, ⁷Kochi Univ., ⁸JAMSTEC, ⁹AIST, ¹⁰LIPI, ¹¹BPPT, ¹²BAKOSURTANAL

A huge ocean-wide tsunami, with average heights of more than 20 meters along the west coast of the northern tip of Sumatra followed the 2004 Sumatra-Andaman earthquake (Mw9.2). Several working hypotheses have been proposed, but the generation mechanism for this tsunami remains unresolved. Most of these hypotheses suggest a possible coseismic slip on splay faults in the outer-arc-high off northwest Sumatra. Among these splay faults, the Middle Thrust (or possibly the Lower Thrust), can best account for features of the Indian Ocean tsunamis observed at regional and ocean-wide distances. To map fault traces and other geological structures that may be contributed by splay fault displacements, we conducted the KY09-09 bathymetry survey offshore northern Sumatra in 2009. The aim of that survey was to identify a fault trace that could be considered a candidate for the Middle Thrust (Hirata et al., 2010).

In early November 2010, we have conducted another high-density survey of the likely source region for the tsunami. This survey consists of a MCS (GI-gun, G=45 cuin and I=105 cuin; true GI-gun mode shooting every 10 sec; a 1,200 m-long, 48 channel solid streamer cable) and a 3.5 kHz Sub-Bottom Profiler (automatic ping intervals depending on water depth). A MNBS bathymetry survey using the SEABEAM 2120, shipboard gravity measurement, and 3-component magnetic measurement have also conducted as well. The survey ship speed was set at averagely 4 knots relative to ground. We designed the acoustic survey lines to cross a series of ridges and troughs parallel to the local trench axis and hence to sample fault traces that are candidates of the Main Thrust, the Lower Thrust, the Middle Thrust, the Upper Thrust in the outer-arc high.

The primary objective of the KH-10-5 cruise are to image detailed deformation structure in the uppermost sediment layers, up to 1 second bsfl in TWT, that are plausibly related to deformation occurred along fault traces. Our final goals are (1) to understand the geological structures in the outer-arc high off northwest Sumatra and their deformation history and (2) to resolve the generation mechanism of the Dec 2004 huge tsunami.

Approximately 480 nautical miles of MCS and SBP data were acquired during the KH-10-5 cruise (Figure 1). During the survey, we produced band-pass filtered, single channel profiles as preliminary results for all the survey lines. We could obtain clear images down to about 1.5 sec (TWT) in the trench fill and a maximum of about 1 sec (TWT) in small troughs in the outer-arc high. In Lines 5 and 6, a kink folding and landward vergent backthrusts were identified near the trench. Many of the small basins on the outer-arc high show deformed sediment layer structures, indicating either folding or faulting. Many SBP profiles also show deformation pattern in the uppermost sediment layers that are consistent with deeper deformation imaged by single-channel data. But some of them seem inconsistent, suggesting a difference in deformation pattern between recent (uppermost) and old (substrata) sedimentation periods. In the region where the Middle thrust is postulated, we found abundant evidences of faulting and folding of the sediment within small basins, along lines 4, 5, 6, 8, 10, 11 and 12. But these results are based on onboard processing and are tentative. We are going to process the MCS data and then interpret detailed geological structure in the near future.

Figure 1

The survey lines (heavy black lines) during the KH10-5 cruise. Main structural features (dashed): WAF, West Andaman Fault; UT, Upper Thrust, ; MT, Middle Thrust; LT, Lower Thrust; M'T, Main Thrust. DF, Deformation Front. UT and LT, are depicted according to Sibuet et al. (2007); MT according to Hirata et al. (2010).

Acknowledgement The KH10-5 cruise was conducted by using the R/V Hakuho-maru. We would like to express our gratitude

to the Captain Takatoshi Seino and the crew for their cooperation and support during the cruise.

キーワード: スマトラ, 海底, 調査, 反射法, サブボトム, 断層
Keywords: sumatra, seafloor, survey, reflection, subbottom, fault