

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



HQR023-P02

会場:コンベンションホール

時間:5月24日 14:00-16:30

エジプト、カルーン湖およびハムラ湖における完新世湖水位変動の復元 The Holocene lake level changes at Lake Qarun and Lake Hamura, Egypt

鹿島 薫^{1*}, 鳴橋 竜太郎²

Kaoru Kashima^{1*}, Ryutaro Naruhashi²

¹九州大学理学研究院地球惑星科学部門, ²東京大学新領域創成科学研究科

¹Faculty of Sciences, Kyushu university, ²University of Tokyo

The joint research project for the inland saline lakes has started with Kyushu university, University of Tokyo and Menoufia University in Egypt. The project is a multidisciplinary research program, focused on the environmental history of the Neolithic and later periods in this region. More in detail, our research is aimed at a high resolution environmental reconstruction in order to make clear the climatic development during the Holocene, and its impact on human societies. We undertook field surveys at inland lakes and Lake Qarun at Fayoum Basin and Lake Hamura at Wadi Natrun in northern Egypt.

The warming and very dry climate could be observed in the data from all our drillings in the early Holocene in Egypt. The three times of fluctuation between arid ? humid environment have occurred after then. The cyclic fluctuation of humidity was also observed in Lake Qarun in Egypt. The lake level of the lake was fell down after the 2200 years BP in the Ptolemaios period, and fluctuated its level with hundreds years intervals.

The above-noted recovery of humidity and its regional variability in the Holocene presumably was one of the major causes for the archaeological events, especially with regard to water availability.

キーワード: 内陸塩性湖沼, 湖水位変動, 完新世, エジプト, 珪藻, ナイル川

Keywords: Inland Saline Lake, Lake Level Changes, Holocene, Egypt, Diatom, River Nile