

# Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



MIS001-03

会場:304

時間:5月25日 14:45-15:00

## 南海トラフ檜野岬海丘において発見された厚さ70mの海底地すべり体の構造 Anatomy of a thick (~70 m) submarine land slide discovered at Kashinozaki Knoll in the Nankai Trough

山口 飛鳥<sup>1\*</sup>, Song Bora<sup>2</sup>, Henry Pierre<sup>3</sup>, 金松 敏也<sup>5</sup>, Kyaw Thu MOE<sup>5</sup>, the Expedition 333 Scientists<sup>5</sup>  
Asuka Yamaguchi<sup>1\*</sup>, Bora Song<sup>2</sup>, Pierre Henry<sup>3</sup>, Toshiya Kanamatsu<sup>5</sup>, MOE Kyaw Thu<sup>5</sup>, Scientists the Expedition 333<sup>5</sup>

<sup>1</sup> 東京大学, <sup>2</sup>Rice University, <sup>3</sup>CEREGE, <sup>4</sup> 海洋研究開発機構, <sup>5</sup>IODP

<sup>1</sup>The University of Tokyo, <sup>2</sup>Rice University, <sup>3</sup>CEREGE, <sup>4</sup>JAMSTEC, <sup>5</sup>IODP

The IODP Expedition 333, a part of NanTroSEIZE project, was operated during December 2010 to January 2011. Three sites including subduction input (Sites C0011 and C0012) and submarine landslide (C0018) were drilled. Site C0012 is located at the top of Kashinozaki Knoll, basement high of the northernmost part of the Philippine Sea Plate in off Kumano region in the Nankai Trough. From 15 to 85 m CSF of Hole C0012C, a steeply-dipping interval (dipping angles of 40-70 degrees to southeastward) probably correspondent with a thick landslide (slump) is discovered. We show the anatomy of this landslide based on the review of shipboard structural, litho- and magneto-stratigraphical, geochemical, and physical property measurement.