

MIS023-03

会場:303

時間:5月22日 09:45-10:15

深部地下圏でのメタン生成と微生物の生態 Unveil the Depth of Methanogenesis in Deep Subsurface and Microbial Ecology

加藤 憲二^{1*}
KATO Kenji^{1*}

¹ 静岡大学理学部
¹ Faculty of Science, Shizuoka University

Bubbles of methane in the ground water coming up from deep subsurface in a sedimentary rock area of Hokkaido, North Island of Japan were shown biological contribution to their origin from stable isotope signature of $\delta\text{-}^{13}\text{C-CH}_4$. Gene sequence of archaeal small subunit rRNA derived from 500 m water showed an existence of various methanogenesis archaeal Genus there. Incubation experiments carried out for the subsample of 500 m ground water revealed that these archaeal community exhibited methanogenesis activity under the in situ or somewhat higher temperature of 30 to 40/45 C with anoxic condition, but not at neither 20 C nor 50 C. Similar incubation applied to the ground water of 140 m deep did not show any methanogenesis under even absolute anoxic condition so far examined for 90 days at both in situ and higher temperature as 16 C and 30 C, respectively, though the stable isotope signature of $\delta\text{-}^{13}\text{C-CH}_4$ in the bubbles obtained from this depth was as low as -57 per mil suggesting methane was biogenic and methanogenesis archaeal genes were found.

キーワード: 地下圏, メタン生成, 微生物の生態
Keywords: Subsurface, Methanogenesis, Microbial Ecology