

MIS023-P05

会場: コンベンションホール

時間: 5月22日 16:15-18:45

## 季節性凍結土壌に生じる一酸化二窒素のアイソトポマー解析 Isotopomer analysis of nitrous oxide produced in a seasonally frozen soil

山崎 哲明<sup>1\*</sup>, 服部 祥平<sup>1</sup>, 柳井 洋介<sup>2</sup>, 豊田 栄<sup>1</sup>, 吉田 尚弘<sup>1</sup>

Tetsuaki Yamazaki<sup>1\*</sup>, Shohei Hattori<sup>1</sup>, Yosuke Yanai<sup>2</sup>, Sakae Toyoda<sup>1</sup>, Naohiro Yoshida<sup>1</sup>

<sup>1</sup> 東京工業大学 総合理工学研究科, <sup>2</sup> 農業・食品産業技術総合研究機構

<sup>1</sup>Tokyo Institute of Technology, <sup>2</sup>NARO

Nitrous oxide (N<sub>2</sub>O) is a potent green house gas of which radiative forcing is about 200 times larger than that of CO<sub>2</sub> and contributes to ozone depletion in the stratosphere. In seasonally frozen soil, it is known that more than 50% of the annual N<sub>2</sub>O emission occurs during spring snowmelt period. In order to investigate the N<sub>2</sub>O production/consumption mechanisms under the soil thawing events, we performed isotopomer analysis of N<sub>2</sub>O at an experimental agriculture field plot in northern Japan. It is suggested that intramolecular <sup>15</sup>N site preference (SP) of N<sub>2</sub>O, that is defined as difference in isotope ratios between central (N<sub>c</sub>) and terminal (N<sub>t</sub>) nitrogen in the N<sub>2</sub>O (NNO) molecule, is a powerful tool to quantitatively analyze the contribution of important production pathways such as nitrification, fungal denitrification, chemo-denitrification and bacterial denitrification together with the presence of N<sub>2</sub>O reduction, in addition to the isotope ratio of N (<sup>15</sup>N) and O (<sup>18</sup>O).

The diffusion chamber method using silicone tube, which is recently applied to gas sampling from each soil depth, is useful method for revealing biological processes in soils. In order to apply this method to isotopomer analysis, diffusion equilibrium of N<sub>2</sub>O (concentration, <sup>15</sup>N, <sup>18</sup>O, SP) was confirmed by laboratory experiment. As a result, no significant difference in concentration and isotope ratios between outside and inside of the silicone tube was detected, suggesting the diffusion method can be applied to in situ soil gas sampling for N<sub>2</sub>O studies.

We, then, investigated the isotopomer signatures of soil N<sub>2</sub>O collected from the seasonally frozen soil under soil thawing events. Isotopomer ratios showed no significant variation during N<sub>2</sub>O production peak in subsoil. Both SP and <sup>15</sup>N of N<sub>2</sub>O were within the range of the reported values for bacterial denitrification, suggesting that N<sub>2</sub>O produced during spring snowmelt period was derived from bacterial denitrification and the contribution of N<sub>2</sub>O reduction was likely to be very low.

キーワード: 一酸化二窒素, アイソトポマー, 凍結土壌, 脱窒

Keywords: Nitrous oxide, Isotopomer, Frozen soil, Denitrification