

MIS027-P01

会場: コンベンションホール

時間: 5月22日 10:30-13:00

## オホーツク海サハリン島沖 Lavrentyev 海底断層南側の新領域で得られたガスハイドレートの特徴 Characteristics of hydrate-bound hydrocarbons retrieved from southern Lavrentyev seabed fault, the Sea of Okhotsk

八久保 晶弘<sup>1\*</sup>, 坂上 寛敏<sup>1</sup>, 南 尚嗣<sup>1</sup>, 山下 聡<sup>1</sup>, 高橋 信夫<sup>1</sup>, 庄子 仁<sup>1</sup>, ジン ヤン<sup>2</sup>, ヴェレシャーギナ オルガ<sup>3</sup>, オブジロフ アナトリー<sup>3</sup>

Akihiro Hachikubo<sup>1\*</sup>, Hirotochi Sakagami<sup>1</sup>, Hirotsugu Minami<sup>1</sup>, Satoshi Yamashita<sup>1</sup>, Nobuo Takahashi<sup>1</sup>, Hitoshi Shoji<sup>1</sup>, Young, K. Jin<sup>2</sup>, Olga Vereshchagina<sup>3</sup>, Anatoly Obzhirov<sup>3</sup>

<sup>1</sup> 北見工業大学, <sup>2</sup> 韓国極地研究所, <sup>3</sup> V.I. イリチェフ太平洋海洋学研究所

<sup>1</sup> Kitami Institute of Technology, <sup>2</sup> Korea Polar Research Institute, <sup>3</sup> Pacific Oceanological Institute, FEB RAS

We report molecular and isotopic compositions of hydrate-bound hydrocarbons in the new seepage sites of offshore Sakhalin Island, the Sea of Okhotsk. More than ten gas seep sites have been discovered since 1990s in the north area of Lavrentyev Fault and hydrate-bearing sediments were recovered (Ginsburg *et al.*, 1993; Hachikubo *et al.*, 2010). These sites often accompany with gas plumes from the sea floor and gas hydrates exist in a shallow sediment layer. Recently, Sakhalin Slope Gas Hydrate (SSGH) project was started from 2007 and we retrieved sediment cores from the southern area of Lavrentyev Fault during the SSGH09 and SSGH10 cruises in 2009-2010. We obtained the samples of hydrate-bound gas and dissolved gas in pore water on board, and we measured molecular and stable isotope compositions of them. Empirical classification of the methane stable isotopes; delta <sup>13</sup>C and delta D according to Whiticar *et al.* (1986) and Schoell (1988) indicated their microbial origin via carbonate reduction. Profiles of methane concentration in the pore water suggested a shallow SMI (sulfate-methane interface). SMI depth was estimated as 30-50cm from the sea floor in the case of hydrate-bearing cores, and around 2m from the sea floor in the case of gas-rich cores. Molecular compositions of hydrate-bound gas were almost the same in both northern and southern areas of Lavrentyev Fault. Stable isotope compositions of hydrate-bound gas were concentrated in the range of -204.6 permil to -196.7 permil for delta D and -66.0 permil to -63.2 permil for delta <sup>13</sup>C in the north area (Hachikubo *et al.*, 2010). On the other hand, both isotopes were more depleted in the south area about 6 permil in <sup>13</sup>C and 7 permil in deuterium, respectively, suggested much more active microbial processes in the shallow sediment. Isotopic difference in delta D between hydrate-bound and dissolved gases was about 5 permil at several sites, indicating that the gas hydrates formed from the current gas in pore water (Hachikubo *et al.*, 2009).

Ginsburg, G.D., V.A. Soloviev, R.E. Cranston, T.D. Lorenson, K.A. Kvenvolden (1993) Gas hydrates from the continental slope, offshore Sakhalin Island, Okhotsk Sea. *Geo-Mar. Lett.*, **13**(1), 41-48, doi:10.1007/BF01204391.

Hachikubo A, O. Khlystov, A. Manakov, M. Kida, A. Krylov, H. Sakagami, H. Minami, N. Takahashi, H. Shoji, G. Kalmychikov, J. Poort (2009) Model of formation of double structure gas hydrates in Lake Baikal based on isotopic data. *Geophys. Res. Lett.*, **36**, L18504, doi:10.1029/2009GL039805.

Hachikubo, A., A. Krylov, H. Sakagami, H. Minami, Y. Nunokawa, H. Shoji, T. Matveeva, Y. K. Jin, A. Obzhirov (2010) Isotopic composition of gas hydrates in subsurface sediments from offshore Sakhalin Island, Sea of Okhotsk. *Geo-Mar. Lett.*, **30**, 313-319, doi:10.1007/s00367-009-0178-y.

Schoell, M. (1988) Multiple origins of methane in the earth. *Chem. Geol.*, **71**, 1-10.

Whiticar, M.J., E. Faber, M. Schoell (1986) Biogenic methane formation in marine and freshwater environments: CO<sub>2</sub> reduction vs. acetate fermentation - Isotope evidence. *Geochim. Cosmochim. Acta*, **50**, 693-709.

キーワード: ガスハイドレート, 安定同位体, オホーツク海

Keywords: gas hydrate, stable isotope, Sea of Okhotsk