

MIS029-04

Room:203

Time:May 23 17:15-17:30

## Observation of aerosol particles at the summit and a base of Mt. Fuji

Kazuhiko Miura<sup>1\*</sup>, Kazuhisa Iinuma<sup>1</sup>, Ai Kajikawa<sup>1</sup>, Syou Suda<sup>1</sup>, Masanori Takeda<sup>1</sup>, Tomoko Hasegawa<sup>1</sup>, Hiroshi Kobayashi<sup>2</sup>, Katsuhiro Nagano<sup>3</sup>, Shungo Kato<sup>4</sup>, Hiroshi Yasuda<sup>5</sup>

<sup>1</sup>F. Sci., Tokyo University of Science, <sup>2</sup>University of Yamanashi, <sup>3</sup>F. Sci.Tec., Tokyo University of Science, <sup>4</sup>Tokyo Metropolitan University, <sup>5</sup>National Institute of Radiological Sci.

Sulfur and organic species originated from ocean make new particles to increase the number of cloud condensation nuclei and change properties of cloud (Charlson et al., 1987). However, in the planetary boundary layer, there are many sea-salt particles that provide surfaces for heterogeneous chemical reactions with sulfur or organic gases. There are a few papers of new particle formation observed in the boundary layer under a high-pressure system (ex. Covert et al., 1996). It suggests that particles are produced in the free atmosphere. As the summit of Mt. Fuji, Japan is usually positioned in the free troposphere, we can measure the variation of aerosol in the free troposphere. Our purpose is studying the new particle formation and particle growing process.

Size distributions from 4.4 nm to 5000 nm in diameter were measured with a scanning mobility particle sizer (SMPS, TSI 3936N25 or 3936L22) and an optical particle counter (OPC, RION KR12 or KC01C), at the summit (3776m, 35.36N, 138.73E) and the base Tarobo (1300 m) in July and August 2006 to 2010. Sample air was dried to lower than 20% with a diffusion dryer.

One topic of the results is a peak measured with SMPS at the summit. This peak of 10 nm appeared at about 11 o'clock on 9th and 10th August 2006 and increased their size during daytime. Volume distributions kept almost constant value during increasing their size. This suggests that this size increasing was due to coagulation and these phenomena happened in the wide area. Moreover, these events did not synchronized at Tarobo. This suggests that these peaks may be the new particle formation by gas to particle conversion in the free troposphere around the summit. The event on 9th was occurred after a passing of a typhoon. It is expected that sulphur and organic species winded up by the typhoon produce new particles in the free troposphere.

The burst of the particles smaller than 20nm in diameter continuing longer than 3 hrs was observed 81 times during four summer seasons (134 days). These events were more observed in the daytime (25) than at night (56). Most of these events did not synchronize at Tarobo.

Origins of air mass of 13 times events in 2010 was estimated with a weather map, a prediction of chemical weather map by CFORS, surface weather elements, and surface gas concentrations. The results showed that the continental and maritime origin were estimated 2 and 5 times, respectively. It is found that the concentration just before the maritime event showed lower than those just before the other events.

### Acknowledgments

This study was partly supported by the Grants-in-Aid for Scientific Research, Category A (Grant No. 17201007), and Category C (Grant No. 22510019), from Japan Society for the Promotion of Science and the Sumitomo Foundation. This work was performed during the period in which the NPO "Valid Utilization of Mt. Fuji Weather Station" maintained the facilities.

### References

- Charlson et al., Nature 326, 655, 1987.  
Covert et al., J. Geophys. Res., 101, 6919, 1996.

Keywords: background aerosol, new particle formation, size distribution, radon, ion, cloud condensation nuclei