

# Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



MSD004-06

Room:301A

Time:May 27 15:30-15:45

## Space-borne imaging observation of the Ionosphere, mesosphere, upper atmosphere, and plasmasphere

Akinori Saito<sup>1\*</sup>, Atsushi Yamazaki<sup>2</sup>, Takeshi Sakanoi<sup>3</sup>, Ichiro Yoshikawa<sup>4</sup>

<sup>1</sup>Graduate School of Science, Kyoto Univer, <sup>2</sup>JAXA/ISAS, <sup>3</sup>PPARC, Tohoku University, <sup>4</sup>The University of Tokyo

ISS-IMAP (Ionosphere, Mesosphere, upper Atmosphere, and Plasmasphere mapping) mission is a space-borne imaging missions to observe the Earth's upper atmosphere, the mesosphere, the ionosphere, the thermosphere and the plasmasphere. It is a scientific mission that installs two imaging instruments on the Exposed Facility of Japanese Experiment Module on the International Space Station, Kibo (EF of ISS-JEM), . The observation is planed to be started in 2011 fiscal year. It will make imaging observation of the Earth's upper atmosphere with visible-light and infrared spectrum imager (VISI) and extra ultraviolet imager (EUVI). The objective of this mission is to clarify the physical mechanism of the following three processes: (1) energy transport process by the atmospheric structures whose horizontal scale is 50-500km in the upper atmosphere (2) process of the plasma transport up to 20,000km altitude (3) effect of the upper atmosphere on the space-borne engineering system. ISS-IMAP will measure the following three parameters in the lower latitude region than 50 degrees: (1) distribution of the atmospheric gravity wave in the mesopause (87km), the ionospheric E-region (95km), and the ionospheric F-region (250km) (2) distribution of the ionized atmosphere in the ionospheric F-region (3) distribution of O<sup>+</sup> and He<sup>+</sup> ions in the ionosphere and plasmasphere. VISI will observe the airglow of 730nm (OH, Alt. 85km), 762nm (O<sub>2</sub>, Alt 95km), 630nm(O, Alt.250km) in the Nadir direction. EUVI will measure the resonant scattering of 30.4nm [He<sup>+</sup>] and 83.4nm [O<sup>+</sup>]. Its field-of-view is 15 degrees, and points the limb of the Earth to observe the vertical distribution of the ions. The scientific objectives and current status of the ISS-IMAP mission will be introduced in the presentation.

Keywords: Ionosphere, Mesosphere, Plasamasphere, Thermosphere, International Space Station, Kibo