

PEM005-04

会場:303

時間:5月26日15:05-15:20

## 太陽風-放射線結合と外帯電子加速 Solar wind-radiation belt coupling via wave-particle interactions

三好 由純 <sup>1</sup>\*, 片岡 龍峰 <sup>2</sup>, 笠原 禎也 <sup>3</sup> Yoshizumi Miyoshi<sup>1</sup>\*, Ryuho Kataoka<sup>2</sup>, Yoshiya Kasahara<sup>3</sup>

1名古屋大学太陽地球環境研究所,2東京工業大学,3金沢大学

<sup>1</sup>STEL, Nagoya University, <sup>2</sup>Tokyo Institute of Technology, <sup>3</sup>Kanazawa University

We investigate the solar wind-radiation belt coupling process, focusing on the large flux enhancement of outer belt electrons associated with high speed coronal hole streams. The flux enhancement tends to occur during the high-speed streams with predominantly southward interplanetary magnetic field (IMF). The IMF dependence can be understood as a result of the internal acceleration of relativistic electrons by wave-particle interactions as follows: The internal acceleration by wave-particle interactions is especially effective when a continuous source of hot electrons is maintained to produce chorus waves for several days. The continuous injection is enhanced during a prolonged period of intense convection and/or substorms associated with southward IMF in high-speed streams. Here we show evidence that the activities of hot electrons, whistler mode chorus waves, and convection/substorms during southward IMF events are clearly different from that during northward IMF events. Based on these results, we propose a model of solar-wind radiation belt coupling in which wave-particle interactions driven by continuous hot electron injections play an important role for the flux enhancement of outer belt electrons.

## キーワード: 放射線帯, 太陽風, 電子加速

Keywords: radiation belts, solar wind, relativistic electron acceleration