

PEM006-P03

会場:コンベンションホール

時間:5月26日 10:30-13:00

全球雷雲活動とOLR及びそれらの太陽活動依存性 Global thunderstorm activities and OLR, and their dependence on solar cycle

高橋 幸弘^{1*}, 佐藤 光輝¹, 宮原 ひろ子², 星野 直哉³, 山下 幸三¹

Yukihiro Takahashi^{1*}, Mitsuteru Sato¹, Hiroko Miyahara², Naoya Hoshino³, Kozo Yamashita¹

¹北海道大学 宇宙理学専攻, ²東京大学 宇宙線研究所, ³東北大学 地球物理学専攻

¹Hokkaido University, ²University of Tokyo, ³Tohoku University

Global ELF observation network, GEON, constructed and operated by Hokkaido University, provides information of each cloud to ground lightning discharge, CG, that is, GPS time, location and energy, as well as Schumann resonance, SR, power, a proxy of global energy proxy of lightning discharge. From the standpoint of the relationship between the effect of solar activity to the climate, lightning activity estimated from the ELF measurement in the frequency range between 1 and 100 Hz and the outgoing longwave Radiation, OLR, an indicator of cloud amount, are examined for their periodicity in the periodic range of about one month. SR power shows about 27 day periodicity in solar maximum years and it becomes elongated toward solar minimum. On the other hand, OLR shows same kind of 27 day periodicity in solar maximum years, but only in the Western Pacific Warm Pool area. Both the spectra of SR and OLR have a peak around 35 days in solar minimum years. The average spectrum of OLR in solar maximum years also shows an enhancement in the range of 50 or 60 days corresponding to the main MJO period. In this paper the relationship between the global lightning distribution based on GEON measurement and OLR are discussed in detail, considering one-month periodicity. Especially synchronization of thunderstorm activity between different longitudes is focused.

キーワード: 太陽活動, 気候, 雷放電, OLR, ELF

Keywords: solar cycle, climate, lightning, thunderstorm, OLR, ELF