

PEM007-10

会場:202

時間:5月27日 11:40-11:55

磁気嵐時のホイッスラー波動の励起および伝搬シミュレーション

Simulation study on the evolution and propagation of whistler waves during storm time

三好 由純^{1*}, V. Jordanova², 松本 洋介¹

Yoshizumi Miyoshi^{1*}, V. Jordanova², Yosuke Matsumoto¹

¹ 名古屋大学太陽地球環境研究所, ² ロスアラモス国立研究所

¹STEL, Nagoya University, ²Los Alamos National Laboratory

In the inner magnetosphere, plasma waves play an essential role for cross-energy coupling processes of different plasma/particle populations via wave-particle interactions. In order to understand the temporal-spatial variation of whistler mode waves in the inner magnetosphere, we simulate the evolution of the electron distribution function and the linear growth rate due to the anisotropies with the RAM electron code [Jordanova and Miyoshi, 2005]. Since the linear growth rate depends on the ratio of the plasma frequency to the gyro-frequency, the intense whistler mode emissions are observed near the plasmapause. In fact, the simulation results for a magnetically active period indicate that whistler mode waves are generated from post-midnight to the dawn-side outside the plasmapause. On the other hand, we do not observe significant wave excitation inside the plasmapause, because hot electrons from the plasma sheet do not access inside the plasmapause. The three-dimensional ray-tracing study indicates that the whistler waves excited outside the plasmapause can propagate into the plasmasphere through multiple reflections at LHR resonance point. Therefore, one of the origin of plasmaspheric hiss is whistler mode waves generated outside the plasmasphere, as suggested by Bortnik et al.[2008].

キーワード: 内部磁気圏, ホイッスラー波動, シミュレーション, エネルギー階層間結合

Keywords: inner magnetosphere, whistler mode waves, simulation, cross-energy coupling