

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



PEM007-P01

会場:コンベンションホール

時間:5月27日 15:00-16:15

ホイッスラーコーラスと高エネルギー電子との波動粒子相互作用 : GEMSIS-RB Wave シミュレーション Wave-particle interaction between whistler chorus and high-energy electrons: GEMSIS-RB Wave simulation

齊藤 慎司^{1*}, 三好 由純¹, 関 華奈子¹
Shinji Saito^{1*}, Yoshizumi Miyoshi¹, Kanako Seki¹

¹ 名古屋大学太陽地球環境研究所

¹STEL, Nagoya University

Scattering process of high energy electrons by whistler chorus in the dipole field is studied to understand electron microbursts associated with whistler chorus by using GEMSIS-RB Wave code. Whistlers are assumed to propagate parallel to the magnetic field, and typical whistler chorus parameters, such as wave frequency, frequency drift rate, and wave amplitude, are assumed based on the spacecraft observations. By using the observed parameters, the GEMSIS-RB Wave code calculates the wave-particle interactions between whistler chorus and the high-energy electrons bouncing along the dipole magnetic field. The code can calculate the precipitation loss of energetic electrons during a few days, considering the micro wave-particle interactions with a few msec. This study focuses on the high energy electron precipitation loss associated with the scattering by whistler chorus with realistic parameters, and applies to understand a physics of high energy electron microburst.

キーワード: 放射線帯, ホイッスラーコーラス, 波動粒子相互作用
Keywords: radiation belt, whistler chorus, wave-particle interaction