

PEM026-09

会場:101

時間:5月24日 16:30-16:45

限られた領域から発生する3次元磁気リコネクションにおけるHall効果の重要性 Hall effects for three-dimensional magnetic reconnection with finite width along the direction of the current

中村 琢磨^{1*}, 中村 るみ², Alexandra Alexandrova², 久保田 康文³
Takuma Nakamura^{1*}, Rumi Nakamura², Alexandra Alexandrova², Yasubumi Kubota³

¹ 宇宙航空研究開発機構・宇宙科学研究所, ² オーストリア宇宙科学研究所, ³ 情報通信研究機構
¹ISAS, JAXA, ²IWF, OEAW, ³NICT

We have performed three-dimensional Hall magnetohydrodynamic (MHD) simulations of magnetic reconnection with finite width along the current direction. Previous MHD simulations have revealed that such localized three-dimensional situations produce the so-called fast reconnection process, in which an Alfvénic fast and narrow jet flows from the localized reconnection region, and the reconnected field lines are strongly piled-up at the head of the jet. Our Hall MHD simulations also confirmed a similar fast reconnection process. On the other hand, our Hall MHD simulations also revealed that the Hall effects critically change the results obtained from MHD simulations regarding the physics of the reconnection region; in the Hall MHD cases, the reconnection region itself broadens in the current and anti-current directions due to the plasma flow related with the current. This means that the location and the size of the reconnection region are unsteady features in Hall MHD regime. In this presentation, we will show the detailed results of our Hall MHD simulations and discuss their implications to the observations in the planetary magnetotail.

キーワード: 磁気リコネクション, ホール効果, 3次元シミュレーション

Keywords: magnetic reconnection, Hall MHD, three-dimensional simulation