

PEM027-P09

会場:コンベンションホール

時間:5月25日10:30-13:00

グローバル MHD シミュレーションによる磁力線固有振動数の日変化の再現 Eigenfrequency of geomagnetic field line in a global MHD simulation

才田 聡子 ¹*, 尾花 由紀 ², 藤田 茂 ³, 田中 高史 ⁴, 山岸 久雄 ⁵ Satoko Saita¹*, Yuki Obana², Shigeru Fujita³, Takashi Tanaka⁴, Hisao Yamagishi⁵

¹ 新領域融合研究センター,² 大阪電気通信大学工学部基礎理工学科,³ 気象庁気象大学校,⁴ 九州大学宙空環境研究セン ター,⁵ 国立極地研究所

¹TRIC, ²Osaka Electro-Communication University, ³Meteorological College, Japan Meteorolog, ⁴SERC, Kyushu University, ⁵National Institute of Polar Research

Shear Alfven waves propagating along the geomagnetic field line form standing waves along filed line. The magnetic field intensity and the plasma mass density affect the velocity of shear Alfven waves. Thus temporal and latitudinal variations in the field-line eigenfrequency are potentially caused by variations in the field-line length, the magnetic field intensity, and the mass density along the field line.

We estimated the eigenfrequency by numerically solving the standing Alfven wave equation along the geomagnetic field in a global MHD simulation. The magnitude of diurnal variation in the eigenfrequency becomes larger according as the latitude increases. In low latitudes, the eigenfrequency agrees with that estimated with empirical magnetospheric model (TS04). However, in high latitudes, the rate of variability estimated in a global MHD simulation is larger than the empirically-based assessment.

In this study, we compared the diurnal variations in the eigenfrequencies observed at the vicinity of Syowa Station, Antarctica, and Dunedin, New Zealand with the estimated eigenfrequencies. We found that the global MHD simulation is adequate to reproduce variations in the eigenfrequency at dawn and dusk. The geomagnetic field lines started are extend toward the night sector.

These distorted field lines are not adequately reproduced by the empirical models.