

# Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



PEM028-02

会場:201B

時間:5月27日 14:30-14:45

## 衝撃波上流域での被加速粒子の空間分布推定 Superdiffusion transport of energetic ions accelerated a shock wave

杉山 徹<sup>1\*</sup>, 塩田 大幸<sup>2</sup>

Tooru Sugiyama<sup>1\*</sup>, Daikou Shiota<sup>2</sup>

<sup>1</sup> 海洋研究開発機構, <sup>2</sup> 理化学研究所

<sup>1</sup>JAMSTEC, <sup>2</sup>RIKEN

We study the transport properties of energetic particles in the upstream region of interplanetary shocks considering the possibility of anomalous diffusion, where the density decay profile has not an exponential profile but a power-law behavior. The ACE spacecraft observations at 1 AU show that the energetic ions with energy of 0.55 ~ 0.76 MeV spatial profiles are well fitted by a power law distribution and we have  $\langle dx^2 \rangle \sim t^a$ , with  $a \sim 1.33 \pm 0.01$ . This implies that particle propagation around a near earth orbit can be intermediate between normal diffusion ( $a = 1$ ) and ballistic motion ( $a = 2$ ) even though the power of the magnetic wave is sufficient large to scatter the particles.

キーワード: 衝撃波, 粒子加速, 拡散現象

Keywords: shock wave, particle acceleration, diffusion process