

PEM031-08

会場:103

時間:5月26日 10:15-10:30

## IMF の南向き excursion 時にカスプで観測された電子密度多重増大現象 Multiple enhancements of the electron density in the cusp for brief southward excursions of IMF

白川 拓也<sup>1</sup>, 田口 聡<sup>1\*</sup>, 細川 敬祐<sup>1</sup>, 小川 泰信<sup>2</sup>

Takuya Shirakawa<sup>1</sup>, Satoshi Taguchi<sup>1\*</sup>, Keisuke Hosokawa<sup>1</sup>, Yasunobu Ogawa<sup>2</sup>

<sup>1</sup> 電気通信大学, <sup>2</sup> 国立極地研究所

<sup>1</sup> Univ. of Electro-Communications, <sup>2</sup> NIPR

A brief southward excursion of IMF is thought to cause a short-period enhancement of the magnetosheath plasma precipitation along the reconnected magnetic field lines, and of the electron density at F2 peak altitudes in the cusp. Our survey for the high time resolution data from EISCAT Svalbard Radar found events in which multiple enhancements of the electron density in the cusp were detected during a brief southward excursion of IMF. One of the clear events shows that three enhancements occurred within about 2 minutes of each other. This time interval is much shorter than the mean time between successive auroral events in the cusp, i.e., 6 min. Simultaneous observations of the plasma precipitation by DMSP spacecraft show that the cusp electron precipitation region has a few spatial boundaries in terms of the energy and energy flux of the precipitating electrons. We consider how these spatial boundaries can be incorporated in the multiple enhancements of the electron density detected by the radar, and present a picture for the phenomenon triggered in the ionospheric cusp by a brief southward excursion of IMF.

キーワード: カスプ, 電子密度, プラズマフロー, 惑星間空間磁場

Keywords: cusp, electron density, plasma flow, IMF