

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



PEM031-P04

会場:コンベンションホール

時間:5月26日 14:00-16:30

オンセット直前に現れるビーズ状オーロラのレーダー観測

Observations of large flow shears around small-scale auroral beads observed at substorm onset

細川 敬祐^{1*}, 門倉 昭², 佐藤 夏雄²

Keisuke Hosokawa^{1*}, Akira Kadokura², Natsuo Sato²

¹ 電気通信大学情報理工学研究科, ² 国立極地研究所

¹University of Electro-Communications, ²National Institute of Polar Research

We present, for the first time, a fine-scale electrodynamic structure behind the auroral beads observed at substorm onset, as inferred from high spatial and temporal resolution measurements of substorm aurora with an all-sky TV camera (ATV) and a coherent HF radar of SuperDARN in Iceland. On the night of September 24, 2006, the ATV observed eastward propagating auroral beads in the initial brightening arc of a substorm just prior to the poleward expansion. During the sequential passage of the beads across the radar beams, the radar detected large velocity flow shears whose magnitude was in excess of 1.5 km/s. The observations suggest that flow shears were located very close to the center of the beads; thus, they corresponded to the horizontal electric fields converging toward the beads, which is consistent with the existence of upward field-aligned currents (FACs) flowing out of the beads. The temporal and spatial resolutions of the current radar measurement were still insufficient for fully resolving the detailed electrodynamic structure behind the fast moving auroral beads. At least, however, we can suggest the existence of localized filamentary FAC structures behind the beads, which played a very important role in connecting the auroral beads in the ionosphere and their source perturbation in the magnetosphere.

キーワード: サブストーム, レーダー観測

Keywords: Substorm, Radar Observations