

PEM031-P11

会場:コンベンションホール

時間:5月26日 14:00-16:30

## 極域電離圏における降下粒子と熱流の関係

## On the relation between particle precipitation and heat flux in the polar ionosphere

小川 泰信<sup>1\*</sup>, 田中 良昌<sup>1</sup>, 元場 哲郎<sup>1</sup>, 宮岡 宏<sup>1</sup>, 門倉 昭<sup>1</sup>

Yasunobu Ogawa<sup>1\*</sup>, Yoshimasa Tanaka<sup>1</sup>, Tetsuo Motoba<sup>1</sup>, Hiroshi Miyaoka<sup>1</sup>, Akira Kadokura<sup>1</sup>

<sup>1</sup> 国立極地研究所

<sup>1</sup>National Institute of Polar Research

We have investigated characteristics of particle precipitation and heat flux based on the European Incoherent Scatter (EISCAT) radar data. The soft particle precipitation is often accompanied by electron heating due to the heat flux from the magnetosphere, and both can be a trigger of ionospheric ion upflow. However, little is known about the relation between the particle precipitation and heat flux (for example, relative locations between them). In this paper, we show their characteristics and relations to auroras using data obtained with EISCAT and optical measurements in November 2008 and February 2011, and discuss the plausible mechanisms determining the relative locations between electron heating and auroras in the polar ionosphere.

キーワード: 極域電離圏, 降下粒子, 熱流, EISCAT

Keywords: polar ionosphere, particle precipitation, heat flux, EISCAT