

PEM031-P18

会場:コンベンションホール

時間:5月26日 14:00-16:30

## アルフベン定在波を用いた内部磁気圏のプラズマ密度分布の推定

Plasma mass density profile of the inner magnetosphere estimated from the frequency of standing Alfvén waves

穂積 裕太<sup>1\*</sup>, 能勢 正仁<sup>2</sup>, 越石 英樹<sup>3</sup>, 松本 晴久<sup>3</sup>, 小原 隆博<sup>3</sup>, 古賀 清一<sup>3</sup>, 五家建夫<sup>4</sup>, 三好 由純<sup>5</sup>

Yuta Hozumi<sup>1\*</sup>, Masahito Nose<sup>2</sup>, Hideki Koshiishi<sup>3</sup>, haruhisa matsumoto<sup>3</sup>, Takahiro Obara<sup>3</sup>, Kiyokazu Koga<sup>3</sup>, Tateo Goka<sup>4</sup>, Yoshizumi Miyoshi<sup>5</sup>

<sup>1</sup> 京大・理・地球惑星科学専攻, <sup>2</sup> 京大・理・地磁気世界資料解析センター, <sup>3</sup> 宇宙航空研究開発機構, <sup>4</sup> 東京都市大学工学部機械システム工学科, <sup>5</sup> 名大・STE 研

<sup>1</sup>SPEL, Kyoto Univ., <sup>2</sup>DACGSM, Kyoto Univ., <sup>3</sup>Japan Aerospace Exploration Agency, <sup>4</sup>Faculty of Engineering, Tokyo City Univ., <sup>5</sup>STEL, Nagoya Univ.

The study of plasma density profile in the inner magnetosphere has been based on electron and H<sup>+</sup> density, and the contribution of heavy ions was mostly neglected. Observations by the retarding ion mass spectrometer (RIMS) onboard Dynamic Explorer (DE-1) in 1981-1982 showed that there is heavy ion enhancement near the plasmapause identified by electron or H<sup>+</sup>. In recent years, these cold heavy ions have attracted attention of researchers as the cause of the ring current ion composition change during magnetic storms. Unfortunately, however, there are no direct observations of the ion composition profile of the plasmasphere, since the DE-1 observations in 1980s. In this study, we intend to estimate the heavy ion composition profile in an indirect method. We determined the plasma mass density profile from the fundamental frequency of the harmonic of toroidal standing Alfvén waves, using magnetic field data obtained by the magnetometer on the TSUBASA satellite (MDS-1, Mission Demonstration Test Satellite-1) which had the geo-stationary transfer orbit. We obtained the plasma mass density profile over L=3-6 from around 40 orbital passes on the noon side to the dusk side. We will show statistical results and discuss the existence of the cold heavy ions.