

PPS002-05

会場:103

時間:5月25日 09:30-09:45

## 火星メタンに関する最新の観測結果 Methane on Mars: Current observations and implication

中川 広務<sup>1\*</sup>, 笠羽 康正<sup>1</sup>, 笠井 康子<sup>2</sup>, 佐川 英夫<sup>2</sup>  
Hiromu Nakagawa<sup>1\*</sup>, Yasumasa Kasaba<sup>1</sup>, YASUKO KASAI<sup>2</sup>, Hideo Sagawa<sup>2</sup>

<sup>1</sup> 東北大学, <sup>2</sup> 独立行政法人通信総合研究所

<sup>1</sup>Tohoku University, <sup>2</sup>NICT

Since the simplest organic molecule, methane (CH<sub>4</sub>), was detected in the Martian atmosphere in 2004, this topic has caused a big interest in the solar system, because its discovery indicates the planet is either biologically or geologically active. Up to now, ground-based and space-born observations have confirmed its presence in the Martian atmosphere (Formisano et al., 2004; Krasnopolsky et al., 2004; Geminale et al., 2008; Mumma et al., 2009; Fonti and Marzo, 2010). However, its reservoirs, release mechanisms, circulation, and sink are still open questions. The high-spectral resolution observation from ground-based telescope indicated that CH<sub>4</sub> showed time variation, and non-uniform distribution, with plume-like features at discrete regions (Mumma et al., 2009). Using the Planetary Fourier Spectrometer (PFS) onboard Mars Express (MEX), Geminale et al. (2011) also reported maps of non-uniformed CH<sub>4</sub> distribution. Although the MEX/PFS and the ground-based observations were generally not collected at the same time nor did they cover identical areas of Mars, the characteristics of the observed CH<sub>4</sub> fields appear to be significantly different. In contrast, photochemistry as currently understood does not produce measurable variations in CH<sub>4</sub> concentrations [Lefevre et al., 2009]. Lefevre et al. (2009) suggested an atmospheric lifetime of less than 200 days is necessary to reproduce the local CH<sub>4</sub> enhancements similar to those recently reported, which implies an unidentified CH<sub>4</sub> loss process that is 600 times faster than predicted by standard photochemistry. In summary, current photochemical models cannot explain the observed presence of CH<sub>4</sub> in the atmosphere of Mars and its reported rapid variations in space and time. The origin and sink of CH<sub>4</sub> on Mars remains a puzzle. In the near future, the observations using heterodyne spectroscopy with an alternative CH<sub>4</sub> band near 7.8  $\mu$ m [Sonnabend et al., 2008] are begun to monitor CH<sub>4</sub> from the ground, and the tunable laser spectrometer measurements is carried out at the Martian surface on the 2011 Mars Science Laboratory (MSL) to identify the isotope of CH<sub>4</sub> [Webster et al., 2011]. Joint orbiter mission being considered by ESA and NASA (JIDT) is also expected to provide the significant scientific progress on this issue by investigating the nature of the trace gas with measurements of temperature and aerosols (Zurek et al., 2011). Here we review the current observations of Martian CH<sub>4</sub> and their implications.

キーワード: 火星, メタン, 生命, 居住適性

Keywords: Mars, Methane, Life, Habitability