

PPS022-16

会場:国際会議室

時間:5月27日 10:15-10:30

準惑星ハウメア衝突族の軌道進化: カイパーベルトの歴史への影響 Dynamical Evolution of Haumea Collisional Family: General Properties and Implications for the Trans-neptunian Belt

ソフィアリカフィカ パトリック^{1*}, Jonathan Horner², 中村 昭子³, 向井 正³
Ptryk Sofia Lykawka^{1*}, Jonathan Horner², Akiko M. Nakamura³, Tadashi Mukai³

¹ 近畿大学 総合社会学部 (天文学分野), ²University of New South Wales, ³ 神戸大学 理学研究科

¹Faculty of Soc. & Nat. Sci., Kinki Univ., ²University of New South Wales, ³Graduate School of Science, Kobe Univ.

Recently, the first collisional family was identified in the trans-neptunian belt, thus revealing the importance of collisions between trans-neptunian objects (TNOs). The family consists of the dwarf planet Haumea and at least nine other ~100 km-sized TNOs. Here, we modeled the long term orbital evolution of an ensemble of family fragments over 4 Gyr. First, we obtained collisional families that reproduced the currently known Haumea's family. In particular, ninety percent of the fragments survived the integrations concentrated in wide regions with the following orbital element ranges: $a \sim 6-12$ AU, $e \sim 0.1-0.15$ and $i \sim 7-10$ deg. Most of the survivors populated the so called classical and detached regions of the trans-neptunian belt, whilst a minor fraction entered the scattered disk reservoir (<1%) or was captured in Neptunian resonances (<10%). In addition, the great majority of fragments displayed negligible long term orbital variations. This implies that the orbital distribution of the intrinsic Haumea's family can constrain the orbital conditions and physics of the collision that created the family billions of years ago. Finally, ~25-40% of the original Haumea family was lost due to planetary ejections or collisions over 4 Gyr.

キーワード: エッジワース・カイパーベルト, 太陽系, ハウメア, 衝突族天体, 海王星, 太陽系外縁天体

Keywords: Edgeworth-Kuiper belt, Solar system, Haumea, Collisional family objects, Neptune, Trans-Neptunian objects (TNOs)