

PPS025-07

会場:101

時間:5月23日 15:45-16:00

超高過飽和ガスから生成するダスト類似物の均質核形成と接合成長 Homogeneous nucleation and coalescence growth of dust analogs in supersaturated vapor

木村 勇気^{1*}, 田中 今日子², 三浦 均¹, 塚本 勝男¹
Yuki Kimura^{1*}, Kyoko Tanaka², Hitoshi Miura¹, Katsuo Tsukamoto¹

¹ 東北大学, ² 北海道大学

¹Tohoku University, ²Hokkaido University

To investigate the homogeneous nucleation and growth process of nanoparticles in vapor phase, interferometric observation was attempted for the first time to the gas evaporation method, which has been a commonly accepted physical production method of nanoparticles. Using the gas evaporation method, fine particles with the size of several nm to ~1 micron are directly produced from the gas phase and recognized nanoparticles have a crystalline habit similar with the bulk crystal even in such tiny particles. When an evaporant is initiated in an inert gas, the evaporated vapor subsequently cools and condenses in the gas atmosphere, i.e., solid grains are obtained via homogeneous nucleation from the vapor phase. Therefore it can be assumed that nucleation occurs far from the equilibrium state, but it is not obvious how far condensation takes place. However, there is no report concerning nucleation and limited study in view of crystal growth for smoke experiment, although significant numbers of smoke experiments have been performed so far. Although there has been reports concerning homogeneous and heterogeneous nucleation from solution phases [1], there has been few reports concerning homogeneous nucleation from a vapor phase in recent years. Homogeneous nucleation rates of droplets were measured as a function of temperature and supersaturation using ethanol and nonane [2, 3]. The nucleation rates were different in several orders of magnitude from the classical nucleation theory.

In preliminary experiment, Tungsten oxide was evaporated by electrical heating of a tungsten wire in a mixture gas of Ar and O₂. WO₃ nanoparticles were formed via homogeneous nucleation and growth during a gas cools following a thermal convection produced by the evaporation source. The degree of supersaturation for nucleation was extremely high, 6.6×10^6 , which was determined from the interferogram. Surface free energy of WO₃ at 1100 K was calculated based on the classical nucleation theory and was 1.38×10^3 erg cm⁻², which is within the reported values. Homogeneously condensed WO₃ nuclei initially maintain their temperature for ~5 ms and then cool down with a rate of $\sim 5 \times 10^4$ K/s. A part of the difference between actual formation rate of produced nanoparticles, which were determined based on a transmission electron microscope, and calculated values based on the semi-phenomenological nucleation theory [4] were well explained if we adopt the idea of coalescence growth.

[1] X. Y. Liu, K. Tsukamoto, M. Sorai, *Langmuir*, 16 (2000) 5499.

[2] J. L. Schmitt, G. W. Adams, R. A. Zalabsky, *J. Chem. Phys.* 77 (1982) 2089.

[3] G. W. Adams, J. L. Schmitt, R. A. Zalabsky, *J. Chem. Phys.* 81 (1984) 5074.

[4] K. K. Tanaka, Katsuyuki Kawamura, Hidekazu Tanaka, and Kiyoshi Nakazawa, *J. Chem. Phys.* 122 (2005) 184514.

キーワード: 核形成, 結晶成長, ナノ粒子, ダスト, 干渉計, その場観察

Keywords: nucleation, crystal growth, nanoparticle, dust, interferometer, in-situ observation