

PPS025-11

会場:101

時間:5月23日 17:00-17:15

高密度分子雲内部におけるラジカル - 分子表面反応による二酸化炭素生成 CO₂ formation through radical-molecule reactions on a solid surface inside dense molecular clouds.

大場 康弘^{1*}, 渡部 直樹¹, 香内 晃¹, 羽馬 哲也¹, ピロネロ バレリオ²
Yasuhiro Oba^{1*}, Naoki Watanabe¹, Akira Kouchi¹, Tetsuya Hama¹, Valerio Pirronello²

¹ 北海道大学低温科学研究所, ² カターニア大学

¹ILTS, Hokkaido Univ., ²Universita di Catania

Solid CO₂ is ubiquitously distributed in icy grain mantles in molecular clouds. Since gas phase reactions cannot explain the observed abundance of CO₂ in those environments and CO is also abundant in the ice, CO₂ is considered to form on the surface of icy grains. It has been experimentally demonstrated that CO₂ is formed in interstellar ice analogues processed by UV, ions, or electrons. Recent astronomical observations found solid CO₂ in dense molecular clouds, where the UV field is weak, implying that there should be additional routes to the formation of CO₂ besides UV photolysis. We performed the experiment on surface reactions of CO with cold OH radicals to investigate a possible CO₂ formation route in dense molecular clouds.

OH radicals were produced by dissociating H₂O molecules in microwave-induced plasma and cooled to 100 K before reaction. CO and OH radicals were continuously codeposited onto an Al substrate at 10-40 K. Reaction products were monitored in-situ by FTIR. We found that the formation of CO₂ occurred at all temperatures investigated. Up to 10 % of CO was converted into CO₂ under the present experimental conditions. We propose that surface reactions of CO with non-energetic OH radicals are potential pathways to the formation of CO₂ in dense molecular clouds.

キーワード: 星間分子雲, 化学進化, 二酸化炭素, ラジカル - 分子反応

Keywords: interstellar molecular clouds, chemical evolution, carbon dioxide, radical-molecule reactions