

SCG008-P03

会場:コンベンションホール

時間:5月26日 14:00-16:30

タイ中部 Hua Hin 地域に分布する花崗岩質岩の Rb-Sr 年代 Rb-Sr ages of granitic rocks from the Hua Hin area, Thailand

柚原 雅樹^{1*}, 亀井淳志², 吉本 紋³, 河上哲生⁴, 上久保 寛⁵, 中野伸彦³, 小山内康人³, Punya Charusiri⁶
Masaki Yuhara^{1*}, Atsushi Kamei², Aya Yoshimoto³, Tetsuo Kawakami⁴, Hiroshi Kamikubo⁵, Nobuhiko Nakano³, Yasuhito Osanai³, Punya Charusiri⁶

¹ 福岡大学, ² 島根大学, ³ 九州大学, ⁴ 京都大学, ⁵ JOGMEC, ⁶ Chulalongkorn 大学

¹ Fukuoka University, ² Shimane University, ³ Kyushu University, ⁴ Kyoto University, ⁵ JOGMEC, ⁶ Chulalongkorn University

The granitic rocks are widely distributed in the Hua Hin area, Thailand. This area is located in the Main-Rang Province (Cobbing et al., 1986) consists mainly of S-type granitic rocks, whose ages range from early Late Triassic to late Early Jurassic (ca. 230-180 Ma) (Sone and Metcalfe, 2008). The petrogenesis of these granitic rocks is explained by partial melting of the Sibumasu crust subducted beneath the Palaeo-Tethys accretionary complex (Sone and Metcalfe, 2008). However, the timing of magma activity and characteristic of source material of granitic rocks in the Hua Hin area are poorly understood. In this paper, we address Rb-Sr isotopic study of granitic rocks from this area.

The granitic rocks in the Hua Hin area are composed of foliated granitic rocks and non-foliated granitic rocks. The formers are the Hub Kapong Gneissic Granite, Hua Hin Gneissic Granite and Pran Buri Gneissic Granite. The Hub Kapong and Hua Hin Gneissic Granites are partly weakly mylonitized K-feldspar porphyritic biotite granite. Some of the Hub Kapong Gneissic Granite includes tourmaline. Parts of Hua Hin Gneissic Granite include garnet and tourmaline. Sillimanite is also contained in some of them (Kawakami et al., 2010). The Pran Buri Gneissic Granite is mylonitic biotite granite. Non-foliated granitic rocks are stock bodies intruded into the Hub Kapong Gneissic Granite, and composed of biotite to two-mica granite. These granitic rocks have peraluminous chemical composition. The Hub Kapong and Hua Hin Gneissic Granites and non-foliated granitic rocks indicate low magnetic susceptibility ($SI=1.3-21.5 \times 10^{-5}$), but the Pran Buri Gneissic Granite indicates high value ($SI=185-455 \times 10^{-5}$) (Yoshimoto et al., 2010).

The Hub Kapong and Hua Hin Gneissic Granites give Rb-Sr whole-rock isochron ages of 202 ± 22 Ma ($SrI=0.7259 \pm 0.0024; N=13$) and 209 ± 14 Ma ($SrI=0.7258 \pm 0.0034; N=10$), respectively. Because of dispersion, errors of age and SrI are slightly big. The dispersion might be due to effects of deformation and metamorphism. The Pran Buri Gneissic Granite does not give significant isochron age. A body of non-foliated granitic rocks gives a Rb-Sr whole-rock isochron ages of 84 ± 13 Ma ($SrI=0.7356 \pm 0.0015; N=4$). These ages indicate timing of magmatic activity in the Hua Hin area, and accord with U-Th-Pb monazite ages (213 Ma (core), 83 Ma (rim)) of metamorphic rocks from this area (Nakano et al., 2010).

Keywords: Rb-Sr whole-rock isochron age, granitic rocks, Hua Hin, Thailand