

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SCG011-11

Room:304

Time:May 25 11:15-11:30

The correlation between lower ionospheric perturbations as seen by VLF/LF propagation and EQs

Masashi Hayakawa^{1*}, Yasushi Kasahara², takashi nakamura², Yasuhide Hobara², A. A. Rozhnoi³, M. Solovieva³, O. A. Molchanov³

¹UEC AWCC, ²UEC, ³Institute of Physics of the Earth

The subionospheric VLF/LF propagation is extensively used to investigate the lower ionospheric perturbation in possible association with earthquakes. An extensive period of data over 7 yr from January 2001 to December 2007 and a combination of different propagation paths in and around Japan are used to examine the statistical correlation between the VLF/LF propagation anomaly (average nighttime amplitude, dispersion, and nighttime fluctuation) and earthquakes with magnitude >6.0 . It is then found that the propagation anomaly exceeding the 2s (standard deviation) criterion indicating the presence of ionospheric perturbation is significantly correlated with earthquakes with shallow depth (<40 km). Finally, the mechanism of seismoionospheric perturbations is discussed.

Keywords: Earthquakes, VLF/LF propagation, correlation study