

SCG059-11

会場:105

時間:5月26日 14:45-15:00

伊豆・小笠原・マリアナ海洋性島弧の高い地殻生成率 Extremely high crustal production rate of the Izu-Ogasawara-Mariana intra-oceanic arc

高橋 成実^{1*}, 巽好幸¹, 小平秀一¹, 三浦誠一¹, 佐藤壮¹, 山下幹也¹, 野徹雄¹, 瀧澤薫¹, 野口直人¹, 海宝由佳¹, 金田義行¹
Narumi Takahashi^{1*}, Yoshiyuki Tatsumi¹, Shuichi Kodaira¹, Seiichi Miura¹, Takeshi Sato¹, Mikiya Yamashita¹, Tetsuo No¹,
Kaoru Takizawa¹, Naoto Noguchi¹, Yuka Kaiho¹, Yoshiyuki Kaneda¹

¹(独) 海洋研究開発機構

¹JAMSTEC

The crustal structures across the Izu-Ogasawara-Mariana arc, obtained by Japan Agency for Earth-Marine Science and Technology (JAMSTEC), provides an estimate of the average crustal production rate since 50 Ma. It has been well established that continental crusts have been created as the middle crust of the Izu-Ogasawara-Mariana arc with P-wave velocity of 6.0-6.5 km/s (Suyehiro et al., 1996; Takahashi et al., 2007); however, the production rate of this 'continent' has been still unknown yet. The crustal volume can be calculated by crustal structures obtained by seismic surveys, however, a part of the crustal materials are transformed into mantle through differentiation of crustal materials (Takahashi et al., 2007). Based on a model of Tatsumi et al. (2008), we estimated volumes of transformed crustal materials and calculated the total volumes of arc materials. As shown by distribution of high velocity lower crust beneath the eastern half of the Shikoku Basin, the arc volcanisms also occurred on oceanic crusts produced by backarc opening in the past and the crust has been overprinted by the arc activities after stop of the backarc opening. We, therefore, identified the eastern end of the original oceanic crust using magnetic lineation pattern (Okino et al., 1994) and removed volumes of the oceanic crust from total ones of arc crustal materials. It is then suggested that the total volume of crustal materials across the Izu-Ogasawara-Mariana arc is over 16,000 cubic kilometers per one kilometer. The total volume is higher in the northern Izu-Ogasawara arc and smaller in the southern part. We assumed the volumes of the Kyusyu Paleo Ridge as the remnant arc, and found that the crustal production rate of the oceanic arc is unexpected high value. In this presentation, we introduce crustal image across the arc and the detailed scenario derived the result.

キーワード: 海洋性島弧, 地殻構造, 島弧進化, 海底地震計, 構造探査

Keywords: oceanic crust, crustal structure, arc growth, OBS, seismic survey