

SCG060-22

会場:302

時間:5月25日 15:30-15:45

東北日本弧における地殻流体の3次元解析のための電磁気観測 Electromagnetic measurements to image geofluid in three-dimensions under NE-Japan arc

小川 康雄^{1*}, 市来 雅啓², 小山 崇夫³, 藤 浩明⁴, 松島 政貴⁵, 神田 径¹, 吹野 浩美⁵, 本蔵 義守⁵, 上嶋 誠³
Yasuo Ogawa^{1*}, Masahiro Ichiki², Takao Koyama³, Hiroaki TOH⁴, Masaki Matsushima⁵, Wataru Kanda¹, Hiromi Fukino⁵,
Yoshimori Honkura⁵, Makoto Uyeshima³

¹ 東京工業大学火山流体研究センター, ² 東北大学 理学研究科, ³ 東京大学地震研究所, ⁴ 京都大学 理学研究科, ⁵ 東京工業大学理工学研究科

¹VFRC, Tokyo Inst. tech, ²Tohoku University, ³University of Tokyo, ⁴Kyoto University, ⁵Tokyo Institute of Technology

Fluids in the crust play an important role in volcanic processes and earthquake generation processes. Electrical resistivity is a geophysical parameter which is sensitive to the existence and connectivity of fluids. Thus, by imaging the resistivity by electrical induction method such as magnetotellurics, we can get important information on the amount, chemical composition and transport properties of fluids.

We have started magnetotelluric measurements in the NE Japan using wideband (0.01s–1000s) and long period (10s–20,000s) measurements. The wideband measurements are focused around the Onikobe caldera and the surrounding regions. In 2009, Thirty magnetotelluric soundings were carried out in and around the Onikobe caldera. We have found the lower crustal conductor with N-S strike directions and its shallower continuation to the Onikobe caldera, which has E-W directions representing the E-W tectonic compression. In 2010, we had 30 more stations to the south of the Naruko volcano, covering 20km x 20km.

In addition to the wideband measurements, we have made long period measurements to image the upper mantle structure. We have 32 long period stations with 20km grid spacing in order to image the upper mantle.

We will show some preliminary results on those on-going measurements.

キーワード: 地殻流体, マグネトテルリック, 電磁, 比抵抗

Keywords: geofluid, magnetotellurics, electromagnetic method, resistivity