

SCG065-10

会場:301B

時間:5月23日 17:00-17:15

マントル深部へ沈み込む大陸地殻の量の推定 Supply rate of subducting continental crust to the deep mantle

市川 浩樹^{1*}, 河合 研志², 亀山 真典¹, 土屋 卓久¹

Hiroki Ichikawa^{1*}, Kenji Kawai², Masanori Kameyama¹, Taku Tsuchiya¹

¹ 愛媛大学地球深部ダイナミクス研究センター, ² 東京工業大学大学院理工学研究科

¹GRC, Ehime University, ²Tokyo Institute of Technology

Subductions of granitic materials by tectonic erosion, sediment-trapped subduction, and direct subduction of immature oceanic arcs are indicated by recent geological and geophysical surveys. Geological studies have estimated the volume of subducted tonalite-trondhjemite-granodiorite (TTG) materials is 7 times larger than the volume of the present continental crust. Such a huge volume of subducted granitic materials have a large influence on the chemical evolution of the Earth, and therefore their destination in the mantle has long been debated.

A recent study indicates the TTG materials are gravitationally stable in the mantle transition zone. Capacity of this layer is large enough to reserve subducted continental crust, which might control the mantle convection to some extent. However, it is unclear how and how much they subduct dynamically through the mantle under the lithosphere.

In this study, we conducted a numerical and an analytical approach on an entanglement of a granitic layer by a slab. Sustainable thickness of the subduction channel and supply rate of the granitic materials to the deep mantle are estimated.

キーワード: 大陸地殻, スラブ

Keywords: Continental crust, slab