

Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SCG065-13

会場:301B

時間:5月23日 17:45-18:00

第2大陸 The Second Continent

河合 研志^{1*}, 山本 伸次¹, 土屋 卓久², 丸山 茂徳¹
Kenji Kawai^{1*}, Shinji Yamamoto¹, Taku Tsuchiya², Shigenori Maruyama¹

¹ 東京工業大学, ² 愛媛大学

¹Tokyo Institute of Technology, ²Geodynamics Research Center, Ehime Unive

Recent progress in our understanding of the consuming plate boundary indicates the ubiquitous occurrence of tectonic erosion of the hanging wall of the continental margin, sediment-trapped subduction, and direct subduction of immature oceanic arcs into deep mantle. Geological studies have estimated the volume of subducted tonalite?trondhjemite?granodiorite (TTG) materials to about seven times the surface total volume of continental crust. To reveal the fate of subducted crusts and how they recycle within the Earth, we studied high-pressure densities and elastic properties of TTG by means of the first principles computation method and compared them to those of peridotite. We found that TTG is gravitationally stable and its seismic velocities are remarkably faster than peridotite in the depth range from 300 to 800 km, especially from 300 to 670 km. We, therefore, propose SiO₂-rich second continents in the mantle transition zone, which used to form the TTG crust on the Earth's surface. Our proposed model may provide reasonable explanations of seismological observations such as the splitting of the 670 km discontinuity and seismic scatterers in the uppermost part of the lower mantle. The difference in seismic velocities between PREM model and experimental results in the lower part of the transition zone can be explained by 25 volumetric% of TTG, which would correspond to about six times the present volume of the continental crust. Formation and dynamics of those second continents would have controlled the Earth's thermal history over geologic time.

キーワード: 花崗岩, 沈み込み, 第2大陸, 構造浸食, 第一原理計算

Keywords: granite, subduction, second continent, tectonic erosion, first-principle calculation