

SIT002-07

会場:301B

時間:5月27日 10:15-10:30

オマーンオフィオライト海洋地殻の熱水変質に伴う微量元素組成の変化 Trace element distributions of the hydrothermal altered oceanic crust in the Oman ophiolite

山岡 香子^{1*}, 石川 剛志², 川幡 穂高¹
Kyoko Yamaoka^{1*}, Tsuyoshi Ishikawa², hodaka kawahata¹

¹ 東京大学大気海洋研究所, ² 海洋研究開発機構高知コア研究所

¹AORI, Univ. of Tokyo, ²Kochi Core Center, JAMSTEC

Hydrothermal alteration processes of oceanic crust at mid-ocean ridges cause significant changes in elemental budget and vertical distribution. Although previous studies have been reported chemical compositions of oceanic crustal rocks from dredged and/or drilled modern seafloor and ophiolite, available depth-successive data is still limited. In this study, concentrations of trace elements were determined for a complete section of oceanic crust in the Oman ophiolite in order to investigate elemental mobilization during hydrothermal alteration. Pillow basalts altered at low temperature (<100°C) were highly enriched in B, As, Rb, Cs, Ba, U, and moderately enriched in Pb, suggesting that these elements were adsorbed onto and/or incorporated into secondary minerals, such as smectite and calcite. Mn and Zn were enriched in the transition zone between pillow lava and sheeted dike complex, and depleted in base of sheeted dike complex. On the other hand, Cu and Pb of the sheeted dikes were generally depleted. Dolerite dikes in gabbro altered at high temperature (>300°C) showed enrichment of U, indicating addition of U to rock during high temperature alteration. In contrast to the previous views that both Li and B are leached from rocks during hydrothermal alteration at high temperatures, the lower oceanic crust altered >300°C (even at >450°C) showed B-enrichment relative to fresh rocks. This suggests that the altered oceanic crust is a large sink of B and source of Li.

キーワード: 海洋地殻, 熱水変質, 微量元素

Keywords: oceanic crust, hydrothermal alteration, trace element