

SIT002-09

会場:301B

時間:5月27日 11:00-11:15

北西太平洋海洋リソスフェア構造から推定された高速拡大軸における能動的マントル上昇

Active mantle upwelling at fast-spreading ridge deduced from seismic images of old oceanic lithosphere in the NW Pacific

小平 秀一^{1*}, 藤江 剛¹, 野口 直人¹, 山下 幹也¹, 佐藤 壮¹, 高橋 努¹, 海宝 由佳¹, 山本 揚二郎¹, 高橋 成実¹, 西浦 大輔¹, 阪口 秀¹

Shuichi Kodaira^{1*}, Gou Fujie¹, Naoto Noguchi¹, Mikiya Yamashita¹, Takeshi Sato¹, Tsutomu Takahashi¹, Yuka Kaiho¹, Yujiro Yamamoto¹, Narumi Takahashi¹, Daisuke Nishiura¹, Hilde Sakaguchi¹

¹ 海洋研究開発機構

¹JAMSTEC

One of long-standing questions about the mantle flow, which governs an accretion process of oceanic lithosphere, at the mid-oceanic ridges has been whether the mantle upwelling is active or passive. Although there are lines of geological and geophysical evidences which support dominantly passive upwelling at the mid-oceanic ridges in a sense of the global plate tectonics, it is also possible that, when decompression melting occurs, low density melt is preserved in the mantle to create local buoyancy which forms mantle convection near the spreading axis. A few study about ophiolite and gravity anomalies in the East Pacific Rise demonstrated a possible active upwelling diapirs at the ridges, but yet no seismological evidence which directly indicates the active upwelling has been observed. Here, from seismic data acquired at the old Pacific plate (120 ? 130 Ma) off the Kuril trench, we show very high P-wave velocity ($V_p = 8.6$ km/s) and strong anisotropy (7 %) in the uppermost mantle immediately below the oceanic crust having lower crustal reflectors (LCRs) dipping toward the paleo-ridge with dominantly uniform spacing and dipping. Similar LCRs have been reported by previous seismic studies in the northwestern and eastern Pacific. Based on geometry and distribution of the LCRs, there has been much debate about an origin of the LCR. For example, thermal and chemical modeling predicted that the LCRs were lithological layering formed by downward and outward flow from an axial magma chamber due to passive upwelling of mantle. On the other hand, based on ophiolite studies, it is proposed that ridge-ward lower crustal fabrics may be formed by a basal shear at the crust due to the active mantle upwelling. Our new observations present the first direct seismological evidence indicating strong basal shear of the oceanic crust due to the active upwelling of mantle at the mid-oceanic ridge.

キーワード: 海洋リソスフェア, 地震探査, 地殻, マントル, 異方向性

Keywords: Oceanic lithosphere, Seismic imaging, Crust, Mantle, Anisotropy