

# Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SIT040-P03

会場:コンベンションホール

時間:5月23日 10:30-13:00

## 波形インバーンによる南太平洋下の最下部マントルS波速度構造推定 Waveform inversion for S-velocity structure in the lowermost mantle beneath the Southern Pacific

長谷川 慶<sup>1\*</sup>, 小西 健介<sup>1</sup>, ゲラー ロバート<sup>1</sup>, 富士 延章<sup>3</sup>, 河合 研志<sup>2</sup>  
Kei Hasegawa<sup>1\*</sup>, Kensuke Konishi<sup>1</sup>, Robert J. Geller<sup>1</sup>, Nobuaki Fuji<sup>3</sup>, Kenji Kawai<sup>2</sup>

<sup>1</sup> 東大地惑, <sup>2</sup> 東工大地惑, <sup>3</sup> トゥールーズ大学 UPS-OMP; CNRS,IRAP

<sup>1</sup>EPS, University of Tokyo, <sup>2</sup>EPS, Tokyo Institute of Technology, <sup>3</sup>Universite de Toulouse,UPS-OMP;CNRS,IRAP

We conduct waveform inversion for the vertical profile of shear velocity in the D'' layer beneath the Southern Pacific. We use the transverse component of relatively long period broadband waveforms (20-200s), obtained from IRIS for earthquakes from 1993 to 2010. We find lower S-velocity relative to PREM in the depth range from 0-150km above the core-mantle boundary (CMB), and higher S-velocity relative to PREM in the depth range from 150-300km above the CMB. This is consistent with a phase transition from perovskite to post-perovskite. The average S-velocity in D'' is the same as or slightly faster than PREM, which is roughly consistent with previous global D'' velocity models.