

SMP045-10

Room:301B

Time:May 24 16:45-17:00

## Neutron powder diffraction at high pressure using compact opposed anvil cell

Takuo Okuchi<sup>1\*</sup>, Yoshiki Ohno<sup>2</sup>, Shigeo Sasaki<sup>2</sup>, Jun Abe<sup>3</sup>, Hiroshi Arima<sup>3</sup>, Takanori Hattori<sup>3</sup>, Asami Sano<sup>3</sup>, Osakabe Toyotaka<sup>3</sup>, Kazuki Komatsu<sup>4</sup>, Hiroyuki Kagi<sup>4</sup>

<sup>1</sup>Inst. Study Earth Inter., Okayama Univ., <sup>2</sup>Faculty of Engineering, Gifu Univ., <sup>3</sup>Japan Atomic Energy Agency, <sup>4</sup>Grad. School of Science, Univ. of Tokyo

Current status of technical developments for time-of-flight powder neutron diffraction at high pressure using compact opposed anvil cells were reported [1,2]. We conducted the diffraction experiments at J-PARC MLF BL19 "TAKUMI". Neutron diffraction patterns up to 8 GPa has been successfully obtained using the compact cell.

References: [1] T. Okuchi et al., J. Phys. Conf. Ser., 215 (2010) 012188; [2] T. Okuchi et al., Rev. High Press. Sci. Tech., 20, 175-178, 2010.

Keywords: Powder neutron diffraction, opposed anvil cell