

## スリランカにおける "arrested charnockite" の二種類の産状 Two modes of occurrence of "arrested charnockite" in Sri Lanka

山崎 由貴子<sup>1\*</sup>, 池田 剛<sup>1</sup>, 本吉 洋一<sup>2</sup>, 廣井 美邦<sup>3</sup>, プレーム バーナード<sup>4</sup>  
Yukiko Yamasaki<sup>1\*</sup>, Takeshi Ikeda<sup>1</sup>, Yoichi Motoyoshi<sup>2</sup>, Yoshikuni Hiroi<sup>3</sup>, Bernard Prame<sup>4</sup>

<sup>1</sup>九州大学・理・地惑, <sup>2</sup>国立極地研究所, <sup>3</sup>千葉大学大学院理学研究科, <sup>4</sup>スリランカ地質調査所

<sup>1</sup>Earth and Planet. Sci., Kyushu Univ., <sup>2</sup>National Institute of Polar Research, <sup>3</sup>Earth Sci., Chiba Univ., <sup>4</sup>Geological Survey of Sri Lanka

Felsic to intermediate lithologies charnockitized incompletely have been called as "arrested charnockite". These types of high-grade metamorphic rocks occur in Sri Lanka, as well as Antarctica and South India that formerly constituted Gondwanaland. We described the mode of occurrence and chemical composition of the constituent minerals of the arrested charnockite from two outcrops in Sri Lanka. In the first outcrop near Kurunegala, charnockite occurs as a number of patches with lenticular to ovoidal shapes. The long axes of the patches are not parallel to the foliation defined in the surrounding felsic gneiss. The boundary between charnockite patch and surrounding gneiss is obscure. Compositional layerings composed of Hbl-rich domain and Hbl-poor domain disappear gradually into charnockite. In the second outcrop near Kandy, charnockite is developed as a vein along a fracture of Grt-Bt felsic gneiss. The boundary is obscure. The compositional layerings and mineral preferred orientation recognized in the host gneiss are preserved in the charnockite domain.

In the charnockite from the first outcrop, orthopyroxene occurs commonly. The orthopyroxene grains are locally replaced by cummingtonite along rim and cleavage. Orthopyroxene appears at inner portions of the boundary of charnockite patches recognized by naked eye. The modal abundance of orthopyroxene increases with increasing distance from the boundary. In contrast, vein-like charnockite from the second outcrop, many symplectites occur while orthopyroxene is not found. The surrounding gneiss also has symplectites locally. The symplectites are composed of various mineral assemblages, such as Amp+Pl+Ilm+Mag+Cpx, Amp+Pl+Ilm+Mag and Pl+Ilm+Mag.

Amphibole can be divided into two based on the chemical composition. One has higher Ca and Al than another. The former occurs in the host gneiss and also in the charnockite that defines gneissosity. The latter amphibole (cummingtonite) is recognized only in the charnockite and either replaces orthopyroxene in the patchy charnockite or consists of symplectite in the vein-like charnockite. The cummingtonite in the symplectite showing similar composition to that replacing orthopyroxene suggests that the symplectites in the vein-like charnockite were originally orthopyroxene which has broken down during retrograde metamorphism.

キーワード: チャーノックイト, スリランカ

Keywords: charnockite, Sri Lanka