

# Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SRD043-P01

会場:コンベンションホール

時間:5月22日 10:30-13:00

## 拓洋第5海山に分布するマンガクラストの地球化学的特徴およびレアメタル資源ポテンシャル評価

### Major and trace elements geochemistry of Co-rich ferromanganese crust from the #5 Takuyo Seamount, northwestern Pacific

得丸 絢加<sup>1\*</sup>, 浦辺 徹郎<sup>1</sup>, 白井朗<sup>2</sup>, 野崎 達生<sup>3</sup>, 鈴木 勝彦<sup>3</sup>, NT09-02 乗船者一同<sup>1</sup>

Ayaka Tokumaru<sup>1\*</sup>, Tetsuro Urabe<sup>1</sup>, Akira Usui<sup>2</sup>, Tatsuo Nozaki<sup>3</sup>, Katsuhiko Suzuki<sup>3</sup>, NT09-02 cruise member<sup>1</sup>

<sup>1</sup> 東京大学大学院, <sup>2</sup> 高知大学, <sup>3</sup> 独立行政法人海洋研究開発機構

<sup>1</sup>The University of Tokyo, <sup>2</sup>Kochi University, <sup>3</sup>JAMSTEC

Large amount of hydrogenetic cobalt-rich ferromanganese (Fe-Mn) crusts are known to occur on the surface of seamounts or ocean plateaus in the Pacific Ocean. The Fe-Mn crust is one of the most promising seafloor mineral resources enriched in not only Cu, Co, and Ni but also REE and PGE. Here we report the geochemical features of the Fe-Mn crusts collected from the #5 Takuyo Seamount, which is located 120 km west from the Minami-torishima Island, northwestern Pacific.

The amounts of trace elements in the crusts are controlled by the MnO<sub>2</sub>(or Fe<sub>2</sub>O<sub>3</sub>) content: Those of Ni, and Cu are roughly proportional to the MnO<sub>2</sub> contents, while REE, especially HREE, show strong positive correlation with the Fe<sub>2</sub>O<sub>3</sub> content. The Mn/Fe ratios decrease toward the surface of the crusts in all samples taken at various water depths between 965m and 2987m. The Fe/Mn ratios of crust samples from the surface part within 3 mm tend to be higher along with the increase of water depths, which is similar to the vertical profile of dissolved oxygen amount in the Pacific Ocean. The crusts are characterized by high total REE contents (La-Lu) varying from 1267 to 2168 ppm (average 1660 ppm). PAAS-normalized REE patterns show flat patterns with conspicuous positive Ce anomalies. We believe that the Fe-Mn crusts are promising alternative as the source of HREE because of the high REE contents and the large ore reserve.

キーワード: マングクラスト, 微量元素, 拓洋第5海山, レアメタル・レアアース, 資源ポテンシャル評価

Keywords: ferromanganese crust, minor elements, #5 Takuyo Seamount, REE, resource evaluation