

# Japan Geoscience Union Meeting 2011

(May 22-27 2011 at Makuhari, Chiba, Japan)

©2011. Japan Geoscience Union. All Rights Reserved.



SRD043-P06

会場:コンベンションホール

時間:5月22日 10:30-13:00

## Geology, mineralization and alteration in Nehbandan Mahor Mine, West Lut Block of Iran Geology, mineralization and alteration in Nehbandan Mahor Mine, West Lut Block of Iran

Habib Biabangard<sup>1\*</sup>

Habib Biabangard<sup>1\*</sup>

<sup>1</sup>Sistsn and Baluchestan University

<sup>1</sup>Sistsn and Baluchestan University

The study area is situated within the Lut Block 130 Km east of Nehbandan. The oldest exposed rocks are cretaceous limestone and conglomerate. Dacite-rhyodacite lava and pyroclastic rocks erupted over oldest rocks. Volcanic activities in Tertiary time consist of andesite, trachyandesite, basalt-andesite, dacitic tuff and rhyolite in composition. Plutonic rocks mainly consist of granite, diorite, granodiorite and monzonite. Volcanic rocks are K-rich calc-alkaline. The pattern of spider diagram in comparison with mantle, they are enriched in Cs, Ba, Rb, and Zr and depleted in Nb, K and Ti. In this area alteration zones are silicified, propylitic, sericitic and argillic. Mineralization associated with volcanic rocks show signs of Ag, Au, Zn, Pb and Cu geochemical exploration anomalies.

キーワード: Nehbandan, Geochemical exploration, Alteration, Mineralization, Mahor, Iran

Keywords: Nehbandan, Geochemical exploration, Alteration, Mineralization, Mahor, Iran