

SSS024-P02

Room:Convention Hall

Time:May 27 14:00-16:30

The variation recorded by waveform images of the F-net at the 2007 Niigataken Chuetsu-oki EQ - 2: Hourly plot

Yoshiki Sue^{1*}

¹No institution affiliation

1. Preface

The F-net Broadband Seismograph Network, which is composed of around 100 STS-1 and -2 seismometers, is the network for monitoring earthquakes. Their records are open to the public through its homepage⁽¹⁾.

The homepage provides waveform images (daily plot) in addition to digital data of the waveforms. The purpose of the waveform images could be to help determination of period for waveform data, but they can also be data of seismicity. Because the daily plot is a GIF format image, and whose size (unit: KB, Kilo byte) relates to total length of the drawn lines, further relates to total amount of vibration of the ground where the seismometer is installed. The mathematical explanation of the so far mentioned is shown below, where the symbol => shall mean existence of relationship between the two in this document.

Size of F-net waveform image (KB, Kilo byte)

=> Total length of the drawn lines

=> Total amount of vibration of the ground

The size of the image is summation of drawn lines for specified period, thus it is different from waveform, which is an instant value. The elements which affect its size are amplitude (velocity), frequency and duration of vibration, and it seems that amplitude and duration of vibration affect the value very much.

2. Analyses

2.1 There are several analyses already conducted^{(2),(3)}.

2.2 2007 Niigataken Chetsu-oki Earthquake

<Method>

Same methods as those for the Daily plots.

<Results>

* Basically same pattern for the Daily plots are observed with the following exceptions.

* The hourly plots show quicker changes.

* Further the hourly plots contain spike noises.

* For the several days before the EQ, both patterns are different, i.e., the Daily plots show increase, while the Hourly plots do no show such changes.

Gratitude

The author thanks to National Research Institute for Earth Science and Disaster Prevention (NIED) for the use of F-net data.

References

(1) F-net home page <http://www.fnet.bosai.go.jp/top.php?LANG=en>.

(2) Yoshiki Sue 2010 Long-period vibration recorded by waveform images of the F-net Broadband Seismograph Network, Part 1 (In Japanese) SSJ Fall meeting abstract D31-12.

(3) Yoshiki Sue 2010 Long-period vibration recorded by waveform images of the F-net Broadband Seismograph Network, Part 2 (In Japanese) SSJ Fall meeting abstract P3-60.

Keywords: earthquake, F-net, Niigataken Chuetsu-oki