

SSS035-28

会場:国際会議室

時間:5月24日 15:00-15:15

## スロースリップの物質的な証拠 A material record of slow slip in the shallow accretionary prism

濱田 洋平<sup>1\*</sup>, 坂口 有人<sup>2</sup>, 山口 飛鳥<sup>1</sup>, 亀田 純<sup>1</sup>, 木村 学<sup>1</sup>

Yohei Hamada<sup>1\*</sup>, Arito Sakaguchi<sup>2</sup>, Asuka Yamaguchi<sup>1</sup>, Jun Kameda<sup>1</sup>, Gaku Kimura<sup>1</sup>

<sup>1</sup> 東京大学, <sup>2</sup> 独立法人海洋研究開発機構

<sup>1</sup>The University of Tokyo, <sup>2</sup>JAMSTEC

Recent geophysical observations have shown a great variation in seismicity in a shallow accretionary prism, from the coseismic slip of a large earthquake such as the 1944 Tonankai earthquake to slow slip of VLF in Nankai Trough. During the Integrated Ocean Drilling Program (IODP) Nankai Trough Seismogenic Zone Experiment (NanTro SEIZE) Stage 1, the frontal thrust and a megasplay fault were drilled and cored. The measurements of vitrinite reflectance geothermometry were performed within these two major slip zones and revealed the occurrence of the temperature anomaly along the faults. We combined the data with core-scale structure and interpolated the vitrinite reflectance ( $R_o$ ) applying Steinman function. The smoothed data showed that broad peaks exist within certified slip zones and the peak positions locate in black narrow zones in each fault. We simulated  $R_o$  distribution using temperature alteration model and a kinetic method of thermal maturation of vitrinite. The results show that the range of the  $R_o$  distribution is expanding relative to the width of heat generate region and the long-term heating ( $\sim 100$  s) is needed for the proceeding of the maturing reaction rim. Therefore, we concluded that the measured  $R_o$  distributions indicate that the slow slip occurred in the shallow accretionary prism.

キーワード: スロースリップ, ビトリナイト反射率, 化学反応速度論

Keywords: slow-slip, vitrinite refraction, kinetics