

SSS035-32

会場:国際会議室

時間:5月24日 16:00-16:15

過去の付加体前縁断層に見られる粘土鉱物の相転移反応 Prograde clay minerals reactions along an ancient frontal thrust

亀田 純^{1*}, 山本 由弦², 濱田 洋平¹, 木村 学¹

Jun Kameda^{1*}, Yuzuru Yamamoto², Yohei Hamada¹, Gaku Kimura¹

¹ 東京大学, ² 海洋研究開発機構

¹The University of Tokyo, ²JAMSTEC

Fossil imbricate thrusts branching from an ancient decollement in the Miura-Boso accretionary prism are exposed in the southern parts of the Miura and Boso peninsulas in central Japan. A clay mineralogical analysis on the fault rocks of one of the imbricate thrust faults, the Shirako fault (Boso Peninsula), revealed a local progress of transformation reactions over intact surrounding rocks; illitization of mixed-layer illite-smectite (I-S), thickening and/or ordering of discrete illite and chlorite packets, and partial degradation of kaolinite. Shape of I(001)-S(001) reflection in the X-ray diffraction pattern together with mean crystallite thickness suggests that illite content in I-S within the fault gouge increases by ~10 % from the host rock samples. Local temperature anomaly possibly caused by frictional high-velocity slips appears a favorable reason to have promoted these prograde reactions in the clay assemblage, rather than factors such as rock deformation or specific fluid composition. A thermal model coupled with kinetic simulation on illitization in I-S suggests that the reaction is facilitated by several repetitions of high-velocity slips with peak temperature >400 °C. Such slips along the Shirako fault might have been potentially tsunamigenic events.

キーワード: 付加体, 摩擦発熱, スメクタイト, イライト

Keywords: accretionary prism, tsunami, smectite, illite