

SVC051-04

会場:301B

時間:5月22日 15:00-15:15

高圧実験に基づく三宅島火山大船戸期のマグマ溜まり Magma chamber of Ofunato Stage, Miyakejima volcano based on high-pressure experiments

潮田 雅司^{1*}, 高橋 栄一¹, 鈴木 敏弘², 浜田 盛久¹

Masashi Ushioda^{1*}, Eiichi Takahashi¹, Toshihiro Suzuki², Morihisa Hamada¹

¹ 東京工業大学地球惑星科学専攻, ²IFREE/JAMSTEC

¹Earth and Planetary Sciences, Tokyo Tech, ²IFREE/JAMSTEC

Miyakejima is an active volcanic island located about 200km south of Tokyo in Izu-Mariana arc. Tsukui et al. (2001) divided the volcanic activity of the last 10000 years into four stages: 10000-7000 (Ofunato Stage), 4000-2500 (Tsubota Stage), 2500 y.B.P to AD1154 (Oyama Stage) since AD1469 (Shinmio Stage). We performed melting experiments of OFS scoria, which is one of the least fractionated Miyakejima basalt, in Ofunato stage. Experiments were performed in the temperature ranges of 1050-1200C at 1.0, 1.5, 2.0, 2.5kbar using IHPV at the Magma Factory, Tokyo Tech. Based on the experimental results and petrology of OFS, magma chamber in Ofunato Stage was reconstructed. The magma chamber was located at 5~6km depth (~1.5kbar) and water-rich (~3 wt.%) basalt magma crystallized olivine and calcic plagioclase (which is the typical phenocryst assemblage throughout Ofunato Stage) at ~1100C under NNO-buffer. Condition of the magma chamber was maintained almost constant for 3000 years. A series of crystallization trends were calculated using MELTS program (Ghiorso and Sack, 1995), and it is found that andesites erupted in Tsubota Stage can be formed by fractional crystallization of OFS basalt at 1.5kbar under NNO. Postulated water content in magma (~0.6 wt.%), however, is much lower than in Ofunato Stage (~3 wt.%). Accordingly it is suggested that magma chamber has been significantly degassed (0.6 wt.% H₂O in magma) during the dormant period (4000~7000 y.B.P)

キーワード: 三宅島, マグマ溜まり, 高温高圧実験, 大船戸期

Keywords: Miyakejima, Magma chamber, High-pressure experiment, Ofunato Stage