

U005-04

会場:国際会議室

時間:5月26日 09:10-09:30

はやぶさ回収試料の初期分析：大学コンソーシアムチームによる初期分析の概要 Preliminary examination of Hayabusa asteroidal samples: Overview by university consortium team

土山 明^{1*}, 海老原 充², 木村 真³, 北島 富美雄⁴, 小嗣 真人⁵, 伊藤 正一⁶, 長尾 敬介⁷, 中村 智樹⁸, 奈良岡 浩⁴, 野口 高明³, 岡崎 隆司⁴, 上杉 健太郎⁵, 上根 真之⁹, 坂本 尚義⁶, トレバナー アイランド¹⁰, スコット サンドフォード¹¹, マイケル ゴレンスキー¹², 藤村 彰夫¹³, 安部 正真¹³, 矢田 達¹³, 向井 利典¹³, 岡田 達明¹³, 石橋 之宏¹³, 白井 慶¹³, 上野 宗孝¹³, 川口 淳一郎¹³, 吉川 真¹³

Akira Tsuchiyama^{1*}, Mitsuru Ebihara², Makoto Kimura³, Fumio Kitajima⁴, Masato Kotsugi⁵, Shoichi Itoh⁶, Keisuke Nagao⁷, Tomoki Nakamura⁸, Hiroshi Naraoka⁴, Takaaki Noguchi³, Ryuji Okazaki⁴, Uesugi Kentro⁵, Masayuki Uesugi⁹, Hisayoshi Yurimoto⁶, Trevor R. Ireland¹⁰, Scott A. Sandford¹¹, Michael E. Zolensky¹², Akio Fujimura¹³, Masanao Abe¹³, Toru Yada¹³, Toshifumi Mukai¹³, Tatsuaki Okada¹³, Yukihiko Ishibashi¹³, Kei Shirai¹³, Munetaka Ueno¹³, Junichiro Kawaguchi¹³, Makoto Yoshikawa¹³

¹ 大阪大学理学研究科, ² 首都大学東京理工学研究科, ³ 茨城大学理学部, ⁴ 九州大学理学院, ⁵ 高輝度光科学研究センター SPring8, ⁶ 北海道大学理学研究院, ⁷ 東京大学理学系研究科, ⁸ 東北大学理学研究科, ⁹ 大阪大学工学研究科, ¹⁰ オーストラリア国立大学, ¹¹ NASA Ames 研究センター, ¹² NASA Johnson 研究センター, ¹³ 宇宙航空研究開発機構

¹ Graduate School of Sci., Osaka Univ., ² Grad. Sch. Sci. Eng., Tokyo Met. Univ., ³ Coll. of Sci. at Ibaraki Univ., ⁴ Graduate School of Sci., Kyushu Univ., ⁵ JASRI, SPring-8, ⁶ Graduate School of Sci., Hokkaido Univ., ⁷ Graduate School of Sci., Univ. of Tokyo, ⁸ Graduate School of Sci., Tohoku Univ., ⁹ Graduate School of Eng., Osaka Univ., ¹⁰ Res. School of Earth Sci., ANU, ¹¹ NASA Ames Research Center, ¹² NASA Johnson Research Center, ¹³ JAXA

The Hayabusa spacecraft arrived at S-type Asteroid 25143 Itokawa in November 2006, and reveal astounding features of the small asteroid [1]. Near-infrared spectral shape indicates that the surface of this body has a material similar to that of LL5 or LL6 chondrites [2]. Many small particles (at least >1500 particles; mostly <10 micron and some larger particles of >~100 micron) were successfully collected from MUSES-C region. Tentative analysis with SEM/EDX at the curation facility of JAXA showed that olivine (Fa28/-4), low-Ca pyroxene (Fs23+/-6), high-Ca pyroxene (Fs12+/-9Wo38+/-6), plagioclase (Ab86+/-7) are identified as major phases with minor amounts of Fe-Ni metal, chromite, Ca phosphate, silica minerals and K-bearing halite [3]. About fifty particles of 30-150 microns were allocated for the preliminary examination (PE) by university consortium team.

The basic goals of PE are as follows. (1) Characterization of the surface material of Itokawa, such as classification, formation age, and formation process and conditions. (2) Understanding of processes on a preexisting parent body of Itokawa and accretion into Itokawa, such as examination of brecciation, degrees of impact, etc. (3) Understanding of interaction with space environment, such as space weathering, and isotopic compositions of oxygen and noble gases of the solar wind. (4) Finding foreign substances fallen onto the surface, such as carbonaceous and organic materials and differentiated materials (granitic materials and halite).

The policies of PE are follows. (1) As the sample is not suffered from terrestrial contamination, we make maximum consideration to avoid terrestrial contamination, and obtain data that cannot be obtained from meteorites, such as space weathering, solar wind isotopic compositions, and organic materials. (2) We have to obtain as much data as possible effectively even from a small amount of samples by systematic analyses from non-destructive to destructive ways with minimum contamination in upstream analysis.

We have seven sub-teams; (1) mineralogy and petrology, (2) 3-D structures, (3) elemental compositions, (4) isotopic and minor elemental compositions, (5) noble gas, (6) carbonaceous matter, and (7) organic compounds. PE flow chart is mainly divided into two parts: mainstream and individual analysis flows for specific purposes. In the mainstream, allocated particles have been already measured by non-destructive analyses at Spring-8 and KEK, Japan. 3-D structures, mineral compositions and elemental compositions will be determined by micro-tomography, XRD and XRF, respectively. Based on the non-destructive information, each particle will be cut for later destructive analyses by TEM, SEM, EPMA and SIMS. If carbon is present in the cross section, organic analyses will be made with X-PEEM/XANES and TOF-SIMS.

The following three individual analysis flows are scheduled. (i) Carbonaceous matter on the surfaces will be examined by micro-Raman/IR/fluorescence. Organic materials will be extracted from the particles and analyzed. After the extraction, the residual solid particles will move to the elemental analysis by INAA. If radioactivity is lower than the threshold, the particles will move to the mainstream. (ii) Noble gas analysis will be made using mass spectrometry without exposing particles into the

air. (iii) Space weathering will be examined with a TEM. The samples were prepared carefully to avoid oxidation of Fe metal nano-grains. Potted butts will move to the mainstream. Any suitable sample for isotopic analysis of solar wind oxygen, such as a large single particle of metallic iron or iron sulfide, has not been found at this moment. Result of each analysis will be presented by CoIs of the sub-team in this conference.

[1] Fujiwara A. et al. (2006) *Science*, 312, 1330-1334. [2] Abe M. et al. *Science*, 312, 1334-1338. [3] Nakamura T. et al. (2011) abstract in 42nd LPSC.

キーワード: はやぶさ計画, 初期分析, コンドライト, 小惑星, イトカワ, サンプルリターン

Keywords: Hayabusa mission, preliminary examination, chondrite, asteroid, Itokawa, sample return